



**USAID**  
FROM THE AMERICAN PEOPLE

# CLIMATE CHANGE IN MALI: EXPECTED IMPACTS ON PESTS AND DISEASES AFFLICTING LIVESTOCK

AUGUST 2014

This report is made possible by the support of the American people through the U.S. Agency for International Development (USAID). The contents are the sole responsibility of Tetra Tech ARD and do not necessarily reflect the views of USAID or the U.S. Government.

**ARCC**



African and Latin American  
Resilience to Climate Change Project

This report was prepared by Mario Younan<sup>1</sup> and Brent M. Simpson<sup>2</sup> through a subcontract to Tetra Tech ARD.

<sup>1</sup>Consultant

<sup>2</sup>Michigan State University, Department of Agriculture, Food and Resource Economics

This publication was produced for the United States Agency for International Development by Tetra Tech ARD, through a Task Order under the Prosperity, Livelihoods, and Conserving Ecosystems (PLACE) Indefinite Quantity Contract Core Task Order (USAID Contract No. AID-EPP-I-00-06-00008, Order Number AID-OAA-TO-11-00064).

**Tetra Tech ARD Contacts:**

**Patricia Caffrey**

Chief of Party

African and Latin American Resilience to Climate Change (ARCC)

Burlington, Vermont

Tel.: 802.658.3890

Patricia.Caffrey@tetratech.com

**Anna Farmer**

Project Manager

Burlington, Vermont

Tel.: 802.658.3890

Anna.Farmer@tetratech.com

# CLIMATE CHANGE IN MALI: EXPECTED IMPACTS ON PESTS AND DISEASES AFFLICTING LIVESTOCK

AFRICAN AND LATIN AMERICAN RESILIENCE TO CLIMATE CHANGE (ARCC)

AUGUST 2014

# TABLE OF CONTENTS

<b>ABOUT THIS SERIES .....</b>	<b>v</b>
<b>INTRODUCTION TO THE TABLES .....</b>	<b>1</b>
<b>CATTLE.....</b>	<b>9</b>
VECTOR-BORNE DISEASES.....	9
BOVINE ANAPLASMOSIS .....	9
BOVINE BABESIOSIS .....	9
LUMPY SKIN DISEASE.....	10
RIFT VALLEY FEVER (RVF).....	11
TROPICAL THEILERIOSIS .....	12
TRYPANOSOMOSIS .....	13
ENVIRONMENTALLY TRANSMITTED DISEASES.....	14
ANTHRAX.....	14
FLUKE (FASCIOLA).....	15
GASTRO-INTESTINAL HELMINTHS.....	16
SALMONELLOSIS .....	17
ANIMAL-ANIMAL TRANSMITTED DISEASES .....	18
BOVINE BRUCELLOSIS .....	18
CONTAGIOUS BOVINE PLEURO-PNEUMONIA (CBPP).....	19
DERMATOPHILOSIS.....	20
FOOT-AND-MOUTH DISEASE (FMD) .....	21
HEMORRHAGIC SEPTICEMIA (HS).....	22
<b>CAMEL .....</b>	<b>23</b>
VECTOR-BORNE DISEASES.....	23
RIFT VALLEY FEVER (RVF).....	23
SURRA.....	24
TRYPANOSOMOSIS .....	24
ENVIRONMENTALLY TRANSMITTED DISEASES.....	26
ANTHRAX.....	26
GASTRO-INTESTINAL HELMINTHS.....	27
SALMONELLOSIS .....	28
ANIMAL-ANIMAL TRANSMITTED DISEASES .....	29
BRUCELLOSIS .....	29
CAMEL POX.....	30
DERMATOPHILOSIS.....	31
HEMORRHAGIC SEPTICEMIA (HS).....	32
ORF (PUSTULAR DERMATITIS) .....	33
FLUKE (FASCIOLA).....	34

<b>SHEEP AND GOATS .....</b>	<b>35</b>
VECTOR-BORNE DISEASES.....	35
BLUETONGUE .....	35
HEARTWATER.....	36
RIFT VALLEY FEVER (RVF).....	37
TRYPANOSOMOSIS .....	38
ENVIRONMENTALLY TRANSMITTED DISEASES.....	39
ANTHRAX.....	39
CLOSTRIDIOSIS .....	40
FLUKE (FASCIOLA).....	41
FOOTROT.....	42
GASTRO-INTESTINAL HELMINTHS.....	43
SALMONELLOSIS .....	44
ANIMAL-ANIMAL TRANSMITTED DISEASES .....	45
CONTAGIOUS CAPRINE PLEURO-PNEUMONIA (CCPP).....	45
ORF (PUSTULAR DERMATITIS) .....	46
OVINE BRUCELLOSIS .....	47
PESTE DES PETITS RUMINANTS (PPR) (A.K.A. GOAT PLAGUE).....	48
SHEEP & GOAT POX .....	49
<b>CHICKENS .....</b>	<b>50</b>
VECTOR-BORNE DISEASES.....	50
FOWL POX .....	50
ENVIRONMENTALLY TRANSMITTED DISEASES.....	51
AVIAN COCCIDIOSIS .....	51
GASTRO-INTESTINAL HELMINTHS.....	52
GUMBORO DISEASE.....	53
SALMONELLOSIS .....	54
ANIMAL-ANIMAL TRANSMITTED DISEASES .....	55
NEWCASTLE DISEASE (ND).....	55
HIGHLY PATHOGENIC AVIAN INFLUENZA (HPAI).....	56
PULLORUM DISEASE .....	57
MAREK DISEASE .....	57
AVIAN INFECTIOUS BRONCHITIS .....	58
AVIAN MYCOPLASMOSIS.....	58
FOWL CHOLERA.....	59
<b>DONKEYS, MULES, AND HORSES .....</b>	<b>60</b>
VECTOR-BORNE DISEASES.....	60
AFRICAN HORSE SICKNESS.....	60
TRYPANOSOMOSIS (INCL. TRYPANOSMA EVANSI).....	61
ENVIRONMENTALLY TRANSMITTED DISEASES.....	62
GASTRO-INTESTINAL HELMINTHS AND LIVER FLUKE .....	62
GLANDERS.....	63

<b>PIGS</b> .....	<b>64</b>
VECTOR-BORNE DISEASES.....	64
AFRICAN SWINE FEVER (ASF) .....	64

# ABOUT THIS SERIES

## **ABOUT THE STUDIES ON CLIMATE CHANGE VULNERABILITY AND ADAPTATION IN WEST AFRICA**

This document is part of a series of studies that the African and Latin American Resilience to Climate Change (ARCC) project produced to address adaptation to climate change in West Africa. Within the ARCC West Africa studies, this document falls in the subseries on Climate Change in Mali. ARCC also has produced a subseries on Climate Change and Water Resources in West Africa, Climate Change and Conflict in West Africa, and Agricultural Adaptation to Climate Change in the Sahel.

## **THE SUBSERIES ON ADAPTATION TO CLIMATE CHANGE IN MALI**

At the request of the United States Agency for International Development (USAID), ARCC undertook the Mali series of studies to increase understanding of the potential impacts of climatic change in rural Mali and identify means to support adaptation to these impacts. Other documents in the Climate Change in Mali series include: Expected Impacts on Pests and Diseases Afflicting Selected Crops, Impact Modeling of Selected Agricultural Adaptive Practices, Climate Vulnerability Mapping, Key Issues in Water Resources, Organizational Survey and Focus Groups on Adaptive Practices, and An Institutional Analysis of l'Agence de l'Environnement et du Développement Durable (AEDD) and l'Agence Nationale de la Météorologie (Mali-Météo).

# INTRODUCTION TO THE TABLES

The tables present an analysis of the potential impact of a changed climate on the most common diseases afflicting cattle, sheep, goats, camels, donkeys, pigs, and chickens in Mali. Information used in these tables was drawn from over 60 peer-reviewed scholarly journals and books related to animal health. The website of the World Organisation for Animal Health (OIE) was also consulted. This analysis does not include information from theses, technical reports, newspapers, mainstream magazines, or proceedings of conferences.

For each livestock disease identified, the disease status under current climate was assessed, including the geographical range of the endemic zone, the rate of disease outbreaks within endemic zones, the mode of pathogen transmission, and the relative economic importance to livestock owners. These served as the baseline for an assessment of the likely change in risk of infection under climate scenarios for the 2025 to 2050 period.

The fact that projections of changes to the climate of the Sahel for this period are currently uncertain informed the scenarios used. There is agreement among climate models that temperatures will increase, although the models differ on the extent and rate of that change. Precipitation for this region of the world is particularly difficult to model and existing projections based on these models also differ on the long-term evolution of annual rainfall amounts. Different models produce divergent outcomes for the region; a limited number project increased annual rainfall. The models also provide little insight regarding potential changes in geographic distribution of precipitation. Most models project a slight increase in annual rainfall in the central Sahel and a decrease in the western Sahel. Modelling efforts for Mali are not more precise.

Some models project that the onset of the rainy season may be delayed and extreme events increase. Such intra-annual patterns play a critical role in the severity of disease impact. Changes in the frequency of floods and drought, for example, may significantly impact prevalence. Unfortunately, on the whole, model projections do not address intra-annual weather patterns with the necessary accuracy, and such intra-annual changes in patterns were not considered in the analysis.

This uncertainty in projections argues for an analysis based upon simplified climate scenarios. Because projections are considered reliable with regard to temperature yet inconclusive with regard to annual rainfall amounts, the analysis considered two scenarios. One assumes warmer climate with increased rainfall. A second also assumes a warmer climate, but with lower rainfall.

As the climate scenarios used were basic, the potential impacts identified are straightforward; they consist of risk values of change in infestation or outbreak levels. A number of unknowns prevent greater precision. These include uncertainty regarding the impact of new climatic conditions on pathogen transmission and the immune status of livestock populations. Other factors less dependent on climate will also change. Farmers and pastoralists will adopt new techniques for managing diseases and likely adopt livestock breeds with different resistance to various pests and diseases. Owners may also move their livestock to new agro-ecological zones that pose a lesser (or greater) risk of infection.

Further, available research to explore these issues is limited, especially regarding environmentally and animal-to-animal transmitted diseases. Little information exists on the sensitivity of specific livestock pathogens to moisture and humidity in a hot tropical environment with high ultraviolet radiation, making it difficult to gauge the severity of response to changes in climate.

The predictions that follow are based on expert opinion. They are not the result of modeling or experimentation. They are intended to highlight potential areas of concern. For greater precision, dedicated research targeting specific geographic zones, livestock species, disease vectors, and pathogens under consideration will be necessary.

## **PRESENTATION OF THE TABLES**

The report contains two tables: a summary table, followed by a more detailed table. The summary table is organized by animals and the current prevalence and impact of each pest or disease afflicting them; the effects are characterized as “very high,” “high,” and “moderate.” It presents, for each pest or disease, a brief description of the likely impact on livestock populations under the two climate scenarios. Impacts are tailored to the species and pest or disease, but generally reported in terms of likely changes in numbers of animals lost, (higher, stable, or lower).

The second table is organized by species in this order: cattle; camel; sheep and goats; chickens; donkeys, mules, and horses; and pigs. For each species, the table presents the following information concerning vector-borne diseases, environmentally transmitted diseases, and animal-to-animal transmitted diseases: a general description, including modes and conditions of transmission, and geographic regions of prevalence. This is followed by notes regarding the relationships between climate, livestock management, land use, and pathogen transmission. A table concludes the profile, describing the current status of the disease and the potential impact of each of the two climate change scenarios. The table also presents possible mitigation strategies.

## VERY HIGH

*Pests and diseases currently of serious prevalence and impact*

AFFECTED SPECIES	PEST OR DISEASE	CLIMATE IMPACT
<b><u>CAMEL</u></b>	<a href="#"><u>Rift Valley Fever (RVF)</u></a>	Hot/Wet – Expanded area at risk & increased risk of outbreaks → Could become major livestock & human disease problem; higher impact than in cattle. Hot/Dry – Reduced area at risk & decreased risk of RVF outbreaks
	<a href="#"><u>Surra</u></a>	Hot/Wet – <b>No change</b> if camel population relocates; in the absence of relocation, Surra would occur more frequently and cause higher losses Hot/Dry – <b>No change</b> ; camel population relocates
<b><u>SHEEP AND GOATS</u></b>	<a href="#"><u>Rift Valley Fever (RVF)</u></a>	Hot/Wet – Expanded area at risk & increased risk of outbreaks → Could become <b>major livestock &amp; human disease problem</b> ; sheep see the <b>highest RVF impact of all livestock</b> ; lower impact in goats Hot/Dry – Reduced area at risk & decreased risk of RVF outbreaks; possibility of <b>new RVF foci</b> in irrigation areas
	<a href="#"><u>Peste Des Petits Ruminants (PPR)</u></a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b> ; more frequent outbreaks possible in highly mobile pastoralist livestock
	<a href="#"><u>Contagious Caprine Pleuro-Pneumonia (CCPP)</u></a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b> ; more frequent outbreaks possible in highly mobile pastoralist livestock
<b><u>CHICKENS</u></b>	<a href="#"><u>Newcastle Disease (ND)</u></a>	Hot/Wet – <b>No change</b> Hot/Dry – More frequent Newcastle Disease outbreaks → <b>increased losses</b>
	<a href="#"><u>Highly Pathogenic avian Influenza (HPAI)</u></a>	Hot/Wet – <b>No major change</b> Hot/Dry – <b>Increased risk</b> because of higher concentration of migratory birds in fewer over-wintering habitats
<b><u>DONKEYS, MULES, AND HORSES</u></b>	<a href="#"><u>African Horse Sickness</u></a>	Hot/Wet – <b>Increased risk</b> of disease introduction Hot/Dry – <b>Reduced risk</b> of disease introduction
<b><u>PIGS</u></b>	<a href="#"><u>African Swine Fever (ASF)</u></a>	Hot/Wet – <b>Increased risk</b> of transmission in free ranging domestic pigs Hot/Dry – Reduced risk of transmission and <b>reduced importance</b> of pig industry (rising feed costs!)

## HIGH/SIGNIFICANT

*Pests and diseases currently of significant prevalence and impact*

AFFECTED SPECIES	PEST OR DISEASE	CLIMATE IMPACT
<b>CATTLE</b>	<a href="#">Rift Valley Fever (RVF)</a>	Hot/Wet – <b>Expanded area at risk</b> & increased risk of outbreaks → Could become major livestock & human disease problem; lower impact than in other livestock Hot/Dry – <b>Reduced area at risk</b> & decreased risk of RVF outbreaks; possibility of new RVF foci in irrigation areas
	<a href="#">Tropical Theileriosis</a>	Hot/Wet – Unlikely to be introduced into Mali Hot/Dry – Unusual livestock movement → Increased <b>risk of introduction</b>
	<a href="#">Trypanosomosis</a>	Hot/Wet – <b>Higher losses</b> because of expanded range of tse-tse vector Hot/Dry – <b>No losses</b> ; tse-tse vector and disease may disappear
	<a href="#">Gastro-Intestinal Helminths</a>	Hot/Wet – Heavier endoparasite burden → <b>Increased losses</b> ; lower impact than in sheep, goats and camel Hot/Dry – Lighter endoparasite burden → <b>Reduced losses</b>
	<a href="#">Lumpy Skin Disease</a>	Hot/Wet – Enhanced opportunities for spread of infected insects → <b>higher production losses</b> Hot/Dry – Reduced opportunities for spread of infected insects → <b>lower production losses</b>
	<a href="#">Foot-and-Mouth Disease (FMD)</a>	Hot/Wet – No change Hot/Dry – <b>No change</b> ; more frequent outbreaks possible in highly mobile pastoralist livestock
	<a href="#">Contagious Bovine Pleuro-Pneumonia (CBPP)</a>	Hot/Wet – No change Hot/Dry – No change
<b>CAMEL</b>	<a href="#">Gastro-Intestinal Helminths</a>	Hot/Wet – Heavier endoparasite burden → <b>Increased production losses</b> ; gains much more importance; higher impact than in cattle Hot/Dry – Lighter endoparasite burden → <b>Reduced production losses</b>
	<a href="#">Orf (Pustular dermatitis)</a>	Hot/Wet – More severe clinical disease → <b>higher losses</b> Hot/Dry – Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)
<b>SHEEP AND GOATS</b>	<a href="#">Camel Pox</a>	Hot/Wet – More severe clinical disease → <b>higher losses</b> Hot/Dry – Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)
	<a href="#">Sheep &amp; Goat Pox</a>	Hot/Wet – More severe clinical disease → <b>higher losses</b>

		<b>losses</b> Hot/Dry – Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)
	<a href="#">Orf (Pustular dermatitis)</a>	Hot/Wet – More severe clinical disease → <b>higher losses</b> Hot/Dry – Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)
	<a href="#">Gastro-Intestinal Helminths</a>	Hot/Wet – Heavier endoparasite burden → <b>Increased production losses</b> ; much higher impact than in cattle – <b>likely to gain major importance in sheep and goats</b> Hot/Dry – Lighter endoparasite burden → <b>Reduced production losses</b>
	<a href="#">Footrot</a>	Hot/Wet – Disease occurs much more frequently → significantly <b>increased production losses</b> Hot/Dry – Disease absent over long periods → significantly <b>reduced production losses</b>
<b>CHICKENS</b>	<a href="#">Avian coccidiosis</a>	Hot/Wet – More frequent disease → <b>higher losses</b> Hot/Dry – Less frequent disease → <b>lower losses</b> (modulating effect of poor body condition in relation to affordability/availability of feed)
	<a href="#">Pullorum disease</a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b>
	<a href="#">Marek Disease</a>	Hot/Wet – Slightly <b>increased risk</b> of outbreaks (increased heat stress) Hot/Dry – Slightly <b>increased risk</b> of outbreaks (increased heat stress)
<b>DONKEYS, MULES, AND HORSES</b>	<a href="#">Trypanosomosis</a>	Hot/Wet – Higher transmission risk in wider area → <b>Increased loss of traction power</b> Hot/Dry – Lower transmission risk for <i>T. evansi</i> ; tse-tse transmitted trypanosomes may disappear → <b>Reduced loss of traction power</b>

## MODERATE

*Pests and diseases currently of moderate prevalence and minor impact*

AFFECTED SPECIES	PEST OR DISEASE	CLIMATE IMPACT
<b>CATTLE</b>	<a href="#">Anthrax</a>	Hot/Wet – Less time spent grazing on infected pastures → <b>Lower losses</b> Hot/Dry – Longer time spent grazing on infected pastures → <b>Higher losses</b>
	<a href="#">Bovine Anaplasmosis</a> & <a href="#">Bovine Babesiosis</a>	Hot/Wet – Expanded area at risk → overall <b>increased losses</b> Hot/Dry – Reduced area at risk → overall <b>reduced losses</b>
	<a href="#">Bovine Brucellosis</a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b>
	<a href="#">Fluke (Fasciola)</a>	Hot/Wet – Less time spent grazing risky pasture → <b>Lower losses</b> Hot/Dry – Longer time spent grazing risky pastures → <b>Higher losses</b>
	<a href="#">Dermatophilosis</a>	Hot/Wet – More common occurrence → <b>Significant increase</b> in production <b>losses</b> Hot/Dry – Less common occurrence → <b>Decrease</b> in production <b>losses</b>
	<a href="#">Salmonellosis</a>	Hot/Wet – Enhanced pathogen survival in micro-environments → Slightly <b>increased losses</b> Hot/Dry – Slightly <b>reduced losses</b> esp. under mobile management system
	<a href="#">Hemorrhagic Septicemia (HS)</a>	Hot/Wet – No change Hot/Dry – <b>No change</b> ; potential risk of more severe outbreaks at longer intervals
<b>CAMEL</b>	<a href="#">Anthrax</a>	Hot/Wet – Less time spent grazing on infected pastures → <b>Lower losses</b> Hot/Dry – Longer time spent grazing on infected pastures → <b>Higher losses</b>
	<a href="#">Salmonellosis</a>	Hot/Wet – Enhanced pathogen survival in micro-environments → Slightly <b>increased losses</b> Hot/Dry – Slightly <b>reduced losses</b> esp. under mobile management system
	<a href="#">Dermatophilosis</a>	Hot/Wet – More common occurrence → <b>Increase</b> in production <b>losses</b> ; camels shift north Hot/Dry – Less common occurrence → <b>Decrease</b> or even absence of production <b>losses</b> ; camels kept in arid areas
	<a href="#">Hemorrhagic Septicemia(HS)</a>	Hot/Wet – <b>More frequent</b> outbreaks in camel regions ( <i>guesswork - no information on the etiology</i> ) Hot/Dry – <b>Less frequent</b> more severe outbreaks ( <i>guesswork - no information on the etiology</i> )

	<a href="#">Brucellosis</a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b>
	<a href="#">Trypanosomosis</a>	Camel keeping is only possible in free area, Hot/Wet – camel population shifts North Hot/Dry – camel population shifts South
	<a href="#">Fluke</a>	Hot/Wet – Less time spent grazing risky pasture → <b>Lower losses</b> Hot/Dry – Longer time spent grazing risky pastures → <b>Higher losses</b>
<b><u>SHEEP AND GOATS</u></b>	<a href="#">Heartwater</a>	Hot/Wet – Endemic over expanded range → overall <b>higher losses</b> Hot/Dry – Smaller endemic area → overall <b>lower losses</b>
	<a href="#">Bluetongue</a>	Hot/Wet – Endemic over expanded range → overall <b>higher losses</b> Hot/Dry – Smaller endemic area → overall <b>lower losses</b>
	<a href="#">Trypanosomosis</a>	Hot/Wet – <b>Higher losses</b> because of expanded range of tse-tse vector; lower impact than in cattle Hot/Dry – <b>No losses</b> ; tse-tse vector and disease may disappear
	<a href="#">Anthrax</a>	Hot/Wet – Less time spent grazing on infected pastures → <b>Lower losses</b> Hot/Dry – Longer time spent grazing on infected pastures → <b>Higher losses</b>
	<a href="#">Clostridiosis</a>	Hot/Wet – Disease occurs more frequently → <b>Higher losses</b> Hot/Dry – Disease occurs less frequently → <b>Lower losses</b>
	<a href="#">Ovine Brucellosis</a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b>
	<a href="#">Fluke (Fasciola)</a>	Hot/Wet – Less time spent grazing risky pasture → <b>Lower losses</b> Hot/Dry – More time spent grazing risky pasture → <b>Higher losses</b>
	<a href="#">Salmonellosis</a>	Hot/Wet – Enhanced pathogen survival in micro-environments → Slightly <b>increased losses</b> Hot/Dry – Slightly <b>reduced losses</b> esp. under mobile management system
<b><u>CHICKENS</u></b>	<a href="#">Fowl Pox</a>	Hot/Wet – Increased risk of outbreaks and <b>higher</b> production <b>losses</b> Hot/Dry – Reduced risk of outbreaks → <b>lower</b> production <b>losses</b>
	<a href="#">Gastro-Intestinal Helminths</a>	Hot/Wet – Heavier endoparasite burden → <b>increased losses</b> Hot/Dry – Lighter endoparasite burden → <b>reduced losses</b>
	<a href="#">Gumboro Disease</a>	Hot/Wet – <b>Stable losses</b> Hot/Dry – <b>Stable losses</b>

	<a href="#">Salmonellosis</a>	Hot/Wet – <b>Higher losses</b> Hot/Dry – <b>Reduced losses</b> (modulating effect of poor body condition in relation to affordability/availability of feed)
	<a href="#">Avian infectious bronchitis</a>	Hot/Wet – More severe outbreaks (heat stress) → <b>higher production losses</b> Hot/Dry – More severe outbreaks (heat stress) → <b>higher production losses</b>
	<a href="#">Avian mycoplasmosis</a>	Hot/Wet – More severe outbreaks (heat stress) → <b>higher production losses</b> Hot/Dry – More severe outbreaks (heat stress) → <b>higher production losses</b>
	<a href="#">Fowl Cholera</a>	Hot/Wet – <b>No change</b> Hot/Dry – <b>No change</b>
<b><u>DONKEYS, MULES, AND HORSES</u></b>	<a href="#">Gastro-Intestinal Helminths and Liver Fluke</a>	Hot/Wet – Higher exposure → <b>Increased loss of traction power</b> Hot/Dry – Lower exposure → <b>Reduced loss of traction power</b>
	<a href="#">Glanders</a>	Hot/Wet – <b>No CC related change</b> , veterinary control measures can prevent introduction into Mali Hot/Dry – <b>No CC related change</b> , veterinary control measures can prevent introduction into Mali

# CATTLE

## VECTOR-BORNE DISEASES

### BOVINE ANAPLASMOSIS<sup>i</sup>

- Intra-erythrocytic bacteria (*Anaplasma marginale*) causing fever, progressive anemia, jaundice, weight-loss, and death; calves possess innate protection up to nine months of age; if exposed early, cattle acquire lasting immunity; susceptibility increases with age, and infection of non-immune adults leads to severe acute disease and death.
- Tick-transmitted, while mechanical transmission by biting flies and by unclean vaccination injection needles also plays a role.
- Absent from semi-arid and arid regions (“Sahel Savanna”), it poses a risk to migratory herds entering infested grazing areas; reported exposure rates in the sub-humid “Sudan Savanna” vary between 4 percent and 51 percent.
- In sub-humid areas with consistent exposure of cattle to *Anaplasma* and endemic stability clinical disease and production losses are mostly absent.

### BOVINE BABESIOSIS<sup>ii</sup>

- Intra-erythrocytic protozoan parasite (*Babesia spp.*) causing fever, intra-vascular hemolysis, hematuria (“Redwater”), progressive anemia, jaundice, weight-loss, and death; innate protection lasts up to 6 months of age; early exposed cattle acquire good lasting immunity.
- Transmitted exclusively by *Boophilus spp.* ticks.
- Absent from semi-arid and arid regions (“Sahel Savanna”), it poses a risk to migratory herds entering infested grazing areas; reported exposure rates in the sub-humid “Sudan Savanna” vary between 0 percent and 80 percent.
- In sub-humid areas with endemic stability immune-suppression (drought, calving stress) can lead to recurrence of clinical disease.

### Climate change influence on vector, pathogen-transmission, livestock management, land use (for Babesiosis and Anaplasmosis)

- Increased humidity enhances reproduction and feeding activity of ticks, resulting in more frequent and more regular exposure of cattle to TBD; increased aridity has the opposite effect
- Hot temperatures shorten survival of ticks under both CC scenarios; under hot & humid conditions, shorter survival rate is offset by a higher reproductive rate of ticks
- Migration from non-infected areas into endemic areas occurs more frequently under arid CC scenario

		<b>Expected Climate Change</b>	
--	--	--------------------------------	--

Current Disease Status in Mali	Climate Change Scenario	Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in sub-humid and riverine zones of Mali; sporadic cases in non-immune cattle	Hot/Wet		Expanded area at risk → overall <b>increased losses</b>	
	Hot/Dry	Reduced area at risk → overall <b>reduced losses</b>		Tick control in wet years

### LUMPY SKIN DISEASE<sup>iii</sup>

- Viral pathogen causing fever, multiple skin nodules, and lesions on the inner surfaces of the mouth and respiratory tract; self-limiting with low mortality but has a prolonged debilitating effect on cattle.
- Transmitted by biting insects, which can be carried by wind over long distances.
- Occurs in periodic epidemics that affect large cattle numbers in vast regions, including in the semi-arid Sahel Savanna.
- Economic importance because of prolonged course leading to temporary emaciation, cessation of lactation, temporary loss of fertility, permanent damage to hides.

#### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Increased humidity results in higher reproduction rates and longer activity periods of biting insects resulting in a growing population of infected biting insects; increased aridity has the opposite effect
- Wind speed may be important for regional spread of infected insects

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Enhanced opportunities for spread of infected insects → <b>higher production losses</b>	Hot/Wet		Enhanced opportunities for spread of infected insects → <b>higher production losses</b>	Improved vaccination cover
	Hot/Dry	Reduced opportunities for spread of infected insects → <b>lower production losses</b>		

## RIFT VALLEY FEVER (RVF)<sup>iv</sup>

- Viral pathogen causing mass abortions, hepatitis, and deaths in newborn calves (10 percent-70 percent mortality), sporadic disease and death in adult cattle; very dangerous zoonosis.
- Transmitted by mosquitoes (*Aedes* & *Culex*), which can be wind-carried over long distances; RVF outbreaks in East Africa triggered by abnormally high rainfall with flooding and mass-reproduction of the mosquito vectors – no clear correlation between RVF epidemics and rainfall in West Africa.
- Occurs in periodic epidemics that affect several countries, including in the semi-arid Sahel Savanna.
- Sudden mass abortions lead to disruption of reproductive and production cycles; complete disruption of trade (slaughter ban); RVF can cause potentially lethal disease in humans exposed to infected livestock (during abortions, at slaughter).

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Abnormally high rainfall with flooding leads to mass-reproduction of mosquitoes and a surge in virus transmission; wind speed can affect long distance spread of infected mosquitoes
- In West Africa, expansion of irrigation agriculture has improved conditions for virus transmission significantly

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Evidence from human cases that virus is present in Mali, no outbreaks reported in livestock	Hot/Wet		Expanded area at risk & increased risk of outbreaks → Could become <b>major livestock &amp; human disease problem; lower impact than in other livestock</b>	Improved vaccination cover
	Hot/Dry	Reduced area at risk & decreased risk of RVF outbreaks; possibility of <b>new RVF foci</b> in irrigation areas		

## TROPICAL THEILERIOSIS<sup>vvi</sup>

- Protozoan parasite (*Theileria annulata*) infecting white and red blood cells and causing fever, respiratory symptoms, immune-suppression, diarrhea, anemia, rapid weight loss, and death; young calves are also susceptible, but cattle in endemic regions acquire immunity; mortality of up to 90 percent in naïve cattle.
- Transmitted by *Hyalomma spp.* ticks, which are adapted to arid conditions.
- Present in parts of semi-arid regions (“Sahel Savanna”) of West Africa and in North Africa, it poses a risk of spread into currently non-infected areas and a high risk of severe clinical disease in cattle herds entering infested areas.
- Local endemic stability in permanently infected areas.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Main vector ticks are well adapted to hot and arid conditions, no clear CC effect; potential tick-vectors are already present in non-infected areas
- More frequent long distance transport and migration of livestock due to increased aridity could potentially spread this infection throughout the Sahel region

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Never reported in Mali	Hot/Wet	Unlikely to be introduced into Mali		
	Hot/Dry		Unusual livestock movement → Increased <b>risk of introduction</b>	Strict control of cross-border cattle movement

## TRYPANOSOMOSIS<sup>vii</sup>

- Protozoan blood parasites (in West Africa esp. *Trypanosoma congolense* and *Trypanosoma vivax*, also *Trypanosoma brucei*) cause intermittent fever, lasting immune-suppression, chronic progressive anemia, progressive weight loss (over several months to more than a year), and eventually death. Cattle breeds vary in susceptibility; trypano-tolerant cattle breeds kept in endemic sub-humid to humid regions are in general not suitable for semi-arid Sahel Savanna conditions because they lack drought resistance and good trekking ability.
- Transmitted by tse-tse flies (*Glossina morsitans* group in open savanna habitats and *Glossina palpalis* group in riverine habitats); in adjacent areas without tse-tse flies, transmission of Trypanosomes by biting flies can play a role (especially for *Trypanosoma vivax*).
- Present in all sub-humid regions (“Sudan Savanna”) of West Africa where tse-tse flies occur.
- Leads to complete loss of productivity in infected cattle; unless tse-tse fly vector is efficiently controlled, cattle-keeping becomes uneconomical due to frequent Trypanosomosis.
- Cattle are more affected than sheep and goats because they are much more likely to be bitten by tse-tse flies and to become infected with Trypanosomes.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Increased humidity improves tree cover (long-term) and enhances range, reproduction and feeding activity of tse-tse leading to more frequent infection of cattle with trypanosomes; increased aridity reduces range, reproduction and feeding activity of tse-tse (*G.morsitans*)
- Local emergence of trypanosome transmission by biting flies possible in areas that become unsuitable for tse-tse
- New, very localized, tse-tse habitats inside irrigation areas could emerge (*G. palpalis*)

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Shrinking range of disease and its vector, only present in southern parts of Mali	Hot/Wet		<b>Higher losses</b> because of expanded range of tse-tse vector	Change to more trypano-tolerant cattle breeds
	Hot/Dry	<b>No losses;</b> tse-tse vector and disease may disappear		

## ENVIRONMENTALLY TRANSMITTED DISEASES

### ANTHRAX<sup>viii</sup>

- Spore-forming bacteria (*Bacillus anthracis*) found in soil and particularly in sediment of ponds and lakes; spores survive in the environment for decades, if not centuries; when ingested during grazing or drinking the spores germinate into bacteria and cause a peracute septicemia that leads to death within hours; important zoonosis.
- When carcasses are opened (e.g., by scavengers) bacteria form spores, which again contaminate the environment; spores can be carried by floods onto clean pastures.
- Sporadic outbreaks occur particularly in dry season when animals graze in dry flood zones and gather at overcrowded contaminated watering ponds.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario increased availability of grazing throughout the year allows livestock to spend shorter time grazing infected flood zones
- Increased aridity would see herds spend longer time grazing in high risk areas

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic on certain pastures, esp. in flood zones	Hot/Wet	Less time spent grazing on infected pastures → <b>Lower losses</b>		
	Hot/Dry		Longer time spent grazing on infected pastures → <b>Higher losses</b>	Improved vaccination cover

## FLUKE (FASCIOLA)<sup>ix</sup>

- Trematode parasites (in Sahel especially *Fasciola gigantica*), adult stages parasitize the bile ducts of the liver in cattle; infective stage causes tissue damage due to migration; adult stage causes chronic loss of red blood cells and blood plasma resulting in persistent wasting disease (= typical clinical picture in cattle).
- Indirect transmission cycle includes environmental phase with developmental parasite stages living in water snails (= intermediate hosts); released infectious larvae encyst on vegetation and re-enter cattle host during grazing on fluke-infested pasture.
- Occurs in localized aquatic habitats (near permanent water and in flood zones), including in irrigation schemes (*Fasciola gigantica* gains new habitats through construction of water reservoirs and dams).
- Chronic fasciolosis leads to loss in condition, loss in carcass value, and persistently low milk yield; cattle on high plane of nutrition can tolerate a certain level of fluke infestation.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario creates more suitable snail habitats that support parasite transmission
- Increased aridity reduces habitat range
- Under humid CC scenario herds depend less on fluke-infested grazing; with increased aridity herds are more likely to graze on fluke infested pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in specific habitats (e.g. around water reservoirs, irrigation areas)	Hot/Wet	Less time spent grazing risky pasture → <b>Lower losses</b>		
	Hot/Dry		Longer time spent grazing risky pastures → <b>Higher losses</b>	Strategic anti-parasite treatment

## GASTRO-INTESTINAL HELMINTHS\*

- Gastro-intestinal parasites. Adult stages live in the stomach or intestine; some species attach to the lining of the stomach or the intestine and suck blood. Pathogenicity varies depending on level of infestation and helminth species; they interfere with digestion and some can cause anemia.
- The transmission cycle includes environmental larval phase that live on pasture vegetation. Infective larvae are only active on pasture under humid conditions, and survive only for a rather limited period on pastures under hot and arid conditions; transmission in the semi-arid areas occurs seasonally during rains.
- High infection rate causes wasting disease and interferes with calf growth (stunting); adult cattle develop good resistance against gastro-intestinal helminths but will still be less productive.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario lifespan and activity of infective helminth larvae on pasture increases, resulting in more efficient transmission; increased aridity reduces efficiency of transmission
- High herd mobility in pastoralist system is a modulating factor; humid CC impact felt more in semi-stationary and farm based systems

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in all cattle keeping regions	Hot/Wet		Heavier endoparasite burden → <b>Increased losses</b>	Anti-parasite treatment timed in relation to expected GI-helminth high risk season
	Hot/Dry	Lighter endoparasite burden → <b>Reduced losses</b>		

## SALMONELLOSIS<sup>xi</sup>

- Bacteria living in the gastro-intestinal tract of humans and animals; more than 2000 salmonella serotypes are known worldwide (including some few host specific types) and cause different clinical diseases, including septicemia, diarrhea, and abortions; mortality is seen mainly in young animals; important zoonosis.
- Fecal-oral transmission occurs via diseased and healthy salmonella carriers that contaminate the environment with their feces; salmonella survives for long periods in water and in wet shaded micro-environments.
- Salmonella is more prevalent under intensive husbandry conditions; transmission in extensive production systems in semi-arid areas occurs mainly via crowded and contaminated night enclosures.
- No stable endemic immunity; in combination with the common presence of healthy carriers this leads to persistent losses, especially in young animals; cumulative production losses are often overlooked.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario results in enhanced pathogen survival and multiplication of pathogens in micro-environments and more frequent exposure of cattle to salmonella; higher aridity would lead to reduced exposure
- Transmission via contaminated overnight enclosures in mobile pastoralist systems more affected by increased aridity (= reduced salmonella exposure) than transmission in housed farm animals and transmission on permanent pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet		Enhanced pathogen survival in micro-environments → Slightly <b>increased losses</b>	Improved vaccination cover
	Hot/Dry	Slightly <b>reduced losses</b> esp. under mobile management system		

## ANIMAL-ANIMAL TRANSMITTED DISEASES

### BOVINE BRUCELLOSIS<sup>xii</sup>

- Bacterial pathogen (*Brucella abortus*) causing abortion in cattle and also leading to persistent infections in adult cattle. It is excreted in the milk of healthy carriers (=persistently infected adult cows); important zoonosis in areas with common raw milk consumption.
- Transmission occurs via the environment (stables, enclosures, paddocks), which is contaminated with *Brucella* during abortion; this route plays a by far greater role than occasional venereal transmission; *Brucella* can survive for a short period in humid shaded micro-environments.
- Unless controlled, *Brucella* is common under intensive and extensive husbandry conditions; transmission in extensive production systems in the semi-arid areas occurs mainly via contaminated enclosures.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario results in longer pathogen survival on pasture, resulting in more efficient transmission; increased aridity leads to less efficient transmission via pasture
- Transmission via contaminated pastures and overnight enclosures in mobile pastoralist systems is more affected by aridity (= less efficient transmission) than transmission in housed farm animals on permanent pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet	<b>No change</b>		Improved vaccination cover
	Hot/Dry	<b>No change</b>		

## CONTAGIOUS BOVINE PLEURO-PNEUMONIA (CBPP)<sup>xiii</sup>

- Bacterial pathogen (*Mycoplasma mycoides*) causing an almost invariably chronic progressive respiratory disease in cattle leading to slow wasting. Acute respiratory disease is rare. Persistent infections in adult cattle can be clinically invisible for long intermittent episodes, but stress triggers flare-up of clinical respiratory disease (= chronic cough).
- Transmission occurs via aerosols containing infectious droplets; requires close contact.
- Unless controlled, CBPP is common under intensive and extensive husbandry conditions; animals gradually lose condition/draft power/trekking ability and have to be slaughtered.
- Interferes with national/regional/international trade in livestock.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid conditions favor pathogen survival in infectious aerosols, allowing for slightly more efficient transmission
- Arid conditions lead to poor nutrition status and low immunity in cattle; this reactivates the pathogen excretion by chronic carrier animals and results in more frequent transmission
- Under arid CC scenario there is also a higher risk of pathogen transmission due to high concentration of livestock and more frequent animal contacts at fewer watering points and on overstocked pastures
- Major impact is to be expected more from veterinary control measures than from CC

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali, ongoing control efforts	Hot/Wet	<b>No change</b>		
	Hot/Dry	<b>No change</b>		Improved vaccination cover

## DERMATOPHILOSIS<sup>xiv</sup>

- Bacterial pathogen (*Dermatophilus congolensis*) causing prolonged skin disease with scabs and sometimes generalized skin lesions in cattle, it is common after prolonged rainfall with intense wetting of the skin. It can occasionally occur in suckling calves in the absence of rainy conditions.
- Transmission occurs via direct contact, mostly seasonal.
- Normally no mortality, but has a debilitating effect. Affected cattle have lower milk yield and may lose their calves. Oxen cannot be used for draught power because skin lesions interfere with the yoke; healing takes several weeks.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario would bring about prolonged intensive rainfall with excessive wetting of the skin, which favors more frequent infection and longer duration and intensity of the disease

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Present in Mali and the region, only important in very wet years	Hot/Wet		More common occurrence → <b>Significant increase in production losses</b>	
	Hot/Dry	Less common occurrence → <b>Decrease in production losses</b>		

## FOOT-AND-MOUTH DISEASE (FMD)<sup>xv</sup>

- Viral pathogen causing acute fever and lesions in the mouth and on the feet; it interferes with feeding and walking. Sporadic deaths in small calves and abortions possible.
- The most contagious of all cattle diseases. Transmission via aerosols containing infectious droplets occurs over distances of many kilometers depending on atmospheric conditions. Entire local/national/regional cattle populations may become infected within a short period, and the virus can also be transmitted via milk and meat.
- Unless controlled, FMD outbreaks are common under intensive and extensive husbandry conditions; it disrupts production cycle and is a severe impediment to trade in livestock, meat and milk; **the most important of all trade-relevant diseases in cattle.**

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid conditions favor survival and long distance transmission (several km) of FMD virus in infectious aerosols
- Under arid conditions transmission in pastoralist herds becomes more likely due to more frequent herd contacts at fewer watering points, on overstocked pastures and along trekking routes (this effect is observed empirically but not statistically confirmed )

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet	No change		
	Hot/Dry	<b>No change</b> ; more frequent outbreaks possible in highly mobile pastoralist livestock		Improved vaccination cover

## HEMORRHAGIC SEPTICEMIA (HS)<sup>xvi</sup>

- HS is caused by bacteria (specific capsular type of *Pasteurella multocida*) and presents as a peracute disease with sudden high fever, acute edema, severe respiratory distress, and hemorrhagic septicemia resulting in death within hours.
- Transmission, via aerosols containing infectious droplets, requires close contact; short-term survival of the pathogen in water and role of contaminated watering ponds is being discussed. In between outbreaks, bacteria are found in the respiratory tract of a small percentage of healthy carrier animals. Climatic stress (= sudden transition from hot and dry to wet and cool conditions at the onset of rainy seasons) coupled with poor body condition of cattle at the end of the dry season triggers local outbreaks (multi-factorial disease).
- Outbreaks occur seasonally, but are very sporadic in nature and do not re-occur every year; potentially high mortality.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario episodes with sudden intensive rainfall (= trigger for outbreaks) occur more regularly, but regular exposure and good pasture also enhance the immune status of cattle
- Arid CC scenario would bring poor nutrition and low immunity in cattle, which makes them prone to more severe HS when exposed to sudden intensive rainfall

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Occurs in Mali (seasonal)	Hot/Wet	No change		
	Hot/Dry	<b>No change</b> ; potential risk of more severe outbreaks at longer intervals		Improved vaccination cover

# CAMEL

## VECTOR-BORNE DISEASES

### RIFT VALLEY FEVER (RVF)<sup>xvii</sup>

- Viral pathogen causing mass abortions (80 percent plus in camels), hepatitis, massive deaths in newborn camel calves, and sporadic disease in older camels; very dangerous zoonosis.
- Transmitted by mosquitoes (*Aedes* & *Culex*), which can be carried by wind over long distances. RVF outbreaks in East Africa triggered by abnormally high rainfall with flooding and mass-reproduction of the mosquito vectors; there is no clear correlation between RVF epidemics and rainfall in West Africa.
- Periodic epidemics affect several countries, including in the semi-arid Sahel Savanna.
- Sudden mass abortions lead to disruption of reproductive and production cycle; camels that lose their calves stop lactating and will only calve (and lactate) again one to two years later → massive disruption of milk production and shortage of milk for home consumption in semi-arid and arid regions. RVF can also cause potentially lethal disease in humans exposed to infected livestock (during abortions, at slaughter, and via consumption of raw milk).

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Abnormally high rainfall with flooding leads to mass-reproduction of mosquitoes and a surge in virus transmission
- Wind speed can affect long distance spread of infected mosquitoes
- In West Africa expansion of irrigation agriculture has improved conditions for virus transmission; this may be less significant for camels, which are not kept in irrigation areas

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Evidence from human cases that virus is present in Mali, no outbreaks reported in livestock	Hot/Wet		Expanded area at risk & increased risk of outbreaks  → Could become <b>major livestock &amp; human disease problem; higher impact than in cattle</b>	Improved vaccination cover
	Hot/Dry	Reduced area at risk & decreased risk of RVF		

		outbreaks		
--	--	-----------	--	--

**SURRA<sup>xviii</sup>**

- Protozoan blood parasite (*Trypanosoma evansi*) causes abortions, intermittent fever, immune-suppression, progressive anemia, chronic weight loss over months (in some cases up to two years), and eventually death; calves are less susceptible than adults.
- Transmitted by biting flies (especially *Tabanids* and *Stomoxys spp.*); disease prevalence is linked to seasonal vector density and activity.
- Present in all camel-keeping regions of Asia and Africa including semi-arid parts (“Sahel Savanna”) of West Africa.
- There is no endemic stability; chronically infected camels become unproductive, can no longer be used for transport, and have to be slaughtered. Long-standing infections are refractory to treatment. Economically, Surra is one of the most important diseases in camels.

**Climate change influence on vector, pathogen-transmission, livestock management, land use**

- Increased humidity prolongs activity period of biting flies resulting in a longer transmission period for *T. evansi*
- Increased aridity shortens the period for efficient *T. evansi* transmission
- Camel herds are very mobile, and under humid CC scenario camel-keeping can shift north; with increased aridity camel- keeping could move south

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in camel-keeping areas of Mali	Hot/Wet	No change provided that camel population can relocate; major surge if camel population cannot move		Avoid areas with high density of biting flies
	Hot/Dry	No change; camel population relocates		

**TRYPANOSOMOSIS<sup>xix</sup>**

- In camels, tse-tse transmitted Trypanosomes (in West Africa especially *Trypanosoma congolense* and *Trypanosoma brucei*) causes an acute disease that leads to death within days to a few weeks; very sporadic as camels are normally not kept in Tse-Tse infested areas.
- It is transmitted by tse-tse flies in open savanna and in riverine habitats.
- Present wherever tse-tse flies are common.

- Under constantly high Trypanosome challenge losses occur very frequently; camel- keeping in tse-tse infested areas is not viable.

**Climate change influence on vector, pathogen-transmission, livestock management, land use**

- Increased humidity improves tree cover (long-term) and enhances range, reproduction, and feeding activity of tse-tse (*Glossina morsitans*)
- Increased aridity can lead to emergence of trypanosome transmission by biting flies in areas that become marginal for Tse-Tse
- Under arid CC scenario, *Glossina morsitans* habitats disappear
- Camel herds are very mobile, and under humid CC scenario camel-keeping can shift north; with increased aridity camel- keeping could move south

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Camels are not kept in tse-tse infested zones	Hot/Wet	No change; camel population relocates		Camel-keeping shifts north
	Hot/Dry	No change; camel population relocates		Camel-keeping can expand into new tse-tse free areas (currently infected by tse-tse)

## ENVIRONMENTALLY TRANSMITTED DISEASES

### ANTHRAX<sup>xx</sup>

- Spore-forming bacteria (*Bacillus anthracis*) found in soil and particularly in sediment of ponds and lakes. Anthrax spores survive in the environment for decades if not for centuries. When ingested during grazing or drinking the spores germinate into bacteria and cause a peracute septicemia that leads to death within hours; important zoonosis.
- When carcasses are opened (e.g., by scavengers), bacteria form spores, which again contaminate the environment; spores can be carried by floods onto clean pastures.
- Sporadic outbreaks occur particularly in dry season when animals graze in dry flood zones and gather at overcrowded contaminated watering ponds; less common in camels as compared to cattle.

### Climate change influence on pathogen-transmission, livestock management, land use

- With humid CC scenario camels spend very short time grazing flood zones
- Increased aridity would see camel herds spend longer time in risky grazing areas

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic on certain pastures, esp. in flood zones	Hot/Wet	Less time spent grazing on infected pastures → <b>Lower losses</b>		
	Hot/Dry		Longer time spent grazing on infected pastures → <b>Higher losses</b>	Improved vaccination cover

## GASTRO-INTESTINAL HELMINTHS<sup>xxi</sup>

- Gastro-intestinal parasites; adult stages live in the stomach or intestine; some species attach to the lining of the stomach or the intestine and suck blood. Pathogenicity varies depending on the level of infestation and helminth species; they interfere with digestion and some can cause anemia.
- Transmission cycle includes environmental larval phase that live on pasture vegetation; infective larvae are only active on pasture under humid conditions, and survive only for a rather limited period on pastures under hot and arid conditions; transmission in semi-arid areas occurs seasonally during rains (*Haemonchus longistipes* is specifically adapted to camels and to arid habitats).
- High infection rate causes wasting disease and interferes with camel calf growth (stunting); unlike cattle, adult camels do not develop sufficient resistance against gastro-intestinal helminths → **potentially high impact on production and body condition** of camels.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario lifespan and activity of infective larvae on pasture increases resulting in more efficient transmission; can have major impact unless high camel herd mobility is maintained
- Increased aridity reduces efficiency of helminth larva transmission

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet		Heavier endoparasite burden → <b>Increased production losses</b> ; higher impact than in cattle	Maintain high herd mobility; time anti-parasite treatment in relation to GI-helminth risk period
	Hot/Dry	Lighter endoparasite burden → <b>Reduced production losses</b>		

## SALMONELLOSIS<sup>xxii</sup>

- Bacteria living in the gastro-intestinal tract of humans and animals; more than 2000 *Salmonella* serotypes are known worldwide (including some few host specific types), these cause different clinical diseases, including septicemia, diarrhea, and abortions; mortality is seen mainly in young animals; high prevalence in camel herds; important zoonosis.
- Fecal-oral transmission occurs via diseased and healthy salmonella carriers that contaminate the environment with their feces; salmonella survive for long periods in water and in wet shaded micro-environments.
- Transmission in semi-arid areas occurs mainly via crowded and contaminated night enclosures.
- Lack of stable endemic immunity and common presence of healthy carriers lead to persistent losses, especially in young camels; despite extensive management, salmonella is very prevalent in camels and a major cause of diarrhea and deaths in camel calves – thus limiting herd growth.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario results in enhanced pathogen survival and multiplication of pathogens in micro-environments (overnight enclosures) and more frequent exposure of camels to salmonella
- Humid CC impact can be somewhat minimized by maintaining high camel herd mobility
- Arid CC would lead to reduced salmonella exposure via contaminated overnight enclosures, esp. in very mobile camel herds

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet		Enhanced pathogen survival in micro-environments → Slightly <b>increased losses</b>	Improved vaccination cover, high mobility
	Hot/Dry	Slightly <b>reduced losses</b> , esp. under mobile management system		

## ANIMAL-ANIMAL TRANSMITTED DISEASES

### BRUCELLOSIS<sup>xxiii</sup>

- Bacterial pathogen (*Brucella abortus* and *Brucella melitensis*) causing abortion in pregnant camels and leading to persistent infections in adults (= healthy carriers); important zoonosis in areas with common raw milk consumption.
- The most important *Brucella* transmission route via the environment is less efficient in arid regions where camels are kept.
- *Brucella* is less common in camels as compared to other livestock species.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid scenario results in better pathogen survival on pasture and more efficient transmission
- Increased aridity would reduce efficiency of transmission on pasture
- Transmission via pasture not very efficient in highly mobile camel herds

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet	<b>No change</b>		Improved vaccination cover
	Hot/Dry	<b>No change</b>		

## CAMEL POX<sup>xxiv</sup>

- Viral pathogen (*Orthopoxvirus cameli*) causing fever, abortions, typical pox lesions on the skin, and lesions on the inner surfaces of the respiratory tract; self-limiting, but secondary respiratory infections can cause deaths in weaned young camels; occasionally a peracute form causing death within 24 hours occurs in adult camels that have not been exposed to pox virus before. Strains can differ in virulence.
- Transmitted by direct contact; under humid conditions biting insects transmit bacteria that super-infect skin lesions, which significantly prolongs the healing process and weakens the camel severely.
- Occurs in periodic epidemics that depend largely on the immune status of local camel herds.
- Economic importance because of loss in young camels and interference with breeding and lactation.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid conditions flies transmit bacteria that super-infect pox skin lesions and wet conditions also aggravate secondary respiratory infections that occur after primary Pox infections
- In arid situations the poor nutritional status and low milk yield of lactating camels results in weaker camel calves that are more likely to undergo prolonged severe pox
- More frequent herd contact at watering points may enhance spread of virulent strains

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali and all camel-keeping regions	Hot/Wet		More severe clinical disease → <b>higher losses</b>	Improved vaccination cover (vaccine currently only available in UAE)
	Hot/Dry	Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)		

## DERMATOPHILOSIS<sup>xxv</sup>

- Bacterial pathogen (*Dermatophilus congolensis*) causing prolonged skin disease with scabs, and sometimes generalized skin lesions, after prolonged rainfall with intense wetting of the skin.
- Transmission is via direct contact; occurs only sporadically in camel-keeping regions after exceptionally strong rainfall.
- Normally no mortality, but has a strong debilitating effect on camels (more pronounced than in cattle).

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario would bring longer periods with active infection and transmission over an expanded range

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Present in Mali and the region, only important in very wet years	Hot/Wet		More common occurrence → <b>Increase</b> in production <b>losses</b> ; camels shift north	Migration into more arid area
	Hot/Dry	Less common occurrence → <b>Decrease</b> or even absence of production <b>losses</b> ; camels kept in arid areas		

## HEMORRHAGIC SEPTICEMIA (HS)<sup>xxvi</sup>

- In camels, a disease clinically similar to HS of cattle occurs in devastating outbreaks; climatic stress (= sudden transition from hot and dry to wet and cool conditions at the onset of rainy seasons) and poor body condition play an etiological role. HS in camels is peracute with sudden edema, respiratory distress, and hemorrhagic septicemia resulting in death within hours.
- There is a lack of information on the responsible pathogen and the transmission route; reportedly spread between herds requires close contact (e.g., at watering points).
- Outbreaks are very sporadic in nature with high mortality.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario sudden intensive rainfall occurs more often, outbreaks could occur more frequently
- Arid conditions with poor nutrition lower the immune status and make camels more prone to severe disease
- Under arid conditions introduction of virulent strains is more likely due to more frequent herd contacts

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Not reported from Mali but likely to be present (no information on the etiology, pure guesswork – extrapolated from observations in different camel keeping regions)	Hot/Wet		<b>More frequent</b> outbreaks in camel regions (guesswork – no information on the etiology)	Research into cause of camel HS
	Hot/Dry	<b>Less frequent</b> more severe outbreaks (guesswork – no information on the etiology)		

## ORF (PUSTULAR DERMATITIS)<sup>xxvii</sup>

- Viral pathogen (*Parapoxvirus ovis*) is endemic in all camel-keeping regions. Lesions are similar to camel pox, but affecting mostly the head, lips, and mouth in young suckling camel calves. Healing is without problems unless calves are weak due to lack of milk; occasionally more severe form seen in weaners and sub-adult camels that have not been exposed to the virus before.
- Transmitted by direct contact; Orf virus in camels is possibly related to and shared with sheep and goats.
- Occurs regularly in relation to calving rhythm of the herd.
- Economic importance because of loss or stunting of camel calves.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under wet and unhygienic conditions, secondary bacterial infections prolong healing of Orf lesions resulting in more severe disease
- In arid situations there can be poor nutritional status and low milk yield of lactating camels resulting in weaker camel calves that suffer higher mortality

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali and all camel keeping regions	Hot/Wet		More severe clinical disease → <b>higher losses</b>	Vaccination using sheep Orf vaccine
	Hot/Dry	Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)		

## FLUKE (FASCIOLA)<sup>xxviii</sup>

- Trematode parasites (in Sahel esp. *Fasciola gigantica*), adult stages parasitize the bile ducts of the liver in camels; infective larval stage causes tissue damage due to mass migration through the tissue of the liver; adult stage causes chronic loss of red blood cells and blood plasma resulting in chronic persistent wasting disease (clinical picture in camels is similar to cattle).
- Indirect transmission cycle includes an environmental phase with developmental parasite stages living in water snails (= intermediate hosts), released infectious larvae encyst on vegetation and re-enter camel host during grazing on fluke infested pasture.
- Occurs in localized aquatic habitats (near permanent water and in flood zones), including in irrigation schemes (*Fasciola gigantica* gains new habitats through construction of water reservoirs and dams); due to camels being kept in arid habitats, infection rates in camels are lower than in other ruminants, but camels kept inside or near irrigation areas like the Niger inland delta in Mali can show high infection rates.
- Chronic Fasciolosis leads to loss in condition, loss in carcass value, and persistently low milk yield.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario creates more suitable snail habitats, facilitating parasite transmission
- Increased aridity reduces habitat range
- Under humid CC scenario herds depend less on fluke infested grazing; with increased aridity herds are more likely to graze on fluke infested pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in specific habitats (e.g. around water reservoirs, irrigation areas); <b>minor importance in camels</b>	Hot/Wet	Less time spent grazing risky pasture → <b>Lower losses</b>		
	Hot/Dry		Longer time spent grazing risky pastures → <b>Higher losses</b>	Strategic anti-parasite treatment

# SHEEP AND GOATS

Compared to cattle, Small Ruminant flocks have shorter gestation intervals and higher birth rates and recover faster after drought. This has already led to over-proportional growth of goat/sheep populations. The result is an ever **increasing relative importance of sheep and goat diseases**.

## VECTOR-BORNE DISEASES

### BLUETONGUE<sup>xxix</sup>

- Arthropod-borne viral pathogen (an “Arbovirus”) affecting particularly sheep; it causes severe inflammation of the nasal and oral cavity and of the coronary band above the hooves; in addition there is strong muscle pain and swelling of the head and neck; extreme variability in clinical manifestation (almost invisible in endemic areas); mortality in sheep 2 percent-30 percent ; very mild in goats.
- Transmitted by midges (*Culicoides spp.*), which can be wind-carried over long distances; bluetongue episodes outside endemic humid to sub-humid areas occur in relation to seasonal high rainfall and strong winds; naïve sheep flocks moving from non-endemic arid into endemic sub-humid areas are at particularly high risk of severe disease.
- Severe clinical form requires a prolonged recovery period and renders sheep flock temporarily unproductive.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Higher rainfall leads to enhanced reproduction and biting activity of midges with an increase in virus transmission
- Wind speed affects long distance spread of infected midges
- Under arid conditions migration of naïve herds into endemically infected areas is more likely to occur

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Never reported as clinical disease problem; virus likely to be endemic in sub-humid parts of Mali	Hot/Wet		Endemic over expanded range → overall <b>higher losses</b>	Improved vaccination cover
	Hot/Dry	Smaller endemic area → overall <b>lower losses</b>		

## HEARTWATER<sup>xxx</sup>

- Bacterial pathogen (*Ehrlichia ruminantium*, formerly called *Cowdria ruminantium*), which causes fever, convulsion, central nervous symptoms (circling, in-coordination), and sudden death in goats and sheep. Kids and lambs possess an age related innate resistance. Heartwater susceptibility is breed related, and regional strains can vary in virulence.
- Transmitted by ticks (*Amblyomma spp.* – this tick genus is very widely distributed over tropical Africa); good seasonal rains result in transient increase in heartwater transmission and prevalence.
- Economically more important in Eastern and Southern Africa than in the Sahel – possibly related to differences in virulence between regional *Ehrlichia ruminantium* strains.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Higher rainfall leads to enhanced reproduction and biting activity of ticks and an increase in pathogen transmission
- Under arid conditions migration of naïve herds into endemically infected areas is more likely to occur

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Present wherever vector ticks occur, including in parts of Mali	Hot/Wet		Endemic over expanded range → overall <b>higher losses</b>	Increased use of resistant breeds
	Hot/Dry	Smaller endemic area → overall <b>lower losses</b>		

## RIFT VALLEY FEVER (RVF)<sup>xxx</sup>

- Viral pathogen causing mass abortions, hemorrhagic hepatitis, and 70 percent-100 percent mortality in newborn lambs and kids, and 10 percent-70 percent mortality in older sheep, but less than 10 percent mortality in goats; very dangerous zoonosis.
- Transmitted by mosquitoes (*Aedes* & *Culex*), which can be wind-carried over long distances. RVF outbreaks in East Africa are triggered by abnormally high rainfall with flooding and mass-reproduction of the mosquito vectors, but no clear correlation has been established between RVF epidemics and high rainfall in West Africa.
- Periodic epidemics affect several countries, including in the semi-arid Sahel Savanna.
- Sudden mass abortions and loss of a complete lamb/kid crop lead to complete disruption of reproductive and production cycles; in addition there is also the disruption of trade (slaughter ban). RVF can cause potentially lethal disease in humans exposed to infected livestock, especially sheep, (during abortions, at slaughter).

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Abnormally high rainfall with flooding leads to mass-reproduction of mosquitoes and a sudden surge in virus transmission
- Wind speed can affect long distance spread of infected mosquitoes
- In West Africa, expansion of irrigation agriculture has improved conditions for virus transmission significantly

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Evidence from human cases that virus is present in Mali, no outbreaks reported in livestock	Hot/Wet		Expanded area at risk & increased risk of outbreaks → Could become <b>major livestock and human disease problem; in sheep highest impact of all livestock!</b>	Improved vaccination cover (new RVF vaccine has just become available)
	Hot/Dry	Reduced area at risk and decreased risk of RVF outbreaks; possibility of <b>new RVF foci</b> in irrigation areas		

## TRYPANOSOMOSIS<sup>xxxii</sup>

- Protozoan blood parasites (in West Africa especially *Trypanosoma congolense* and *Trypanosoma vivax*, also *Trypanosoma brucei*) cause intermittent fever, lasting immune-suppression, chronic progressive anemia, progressive weight loss, and eventually death. Sheep breeds vary in susceptibility; trypano-tolerant sheep breeds kept in endemic sub-humid to humid regions are in general not suitable for semi-arid Sahel Savanna conditions because they lack drought resistance and good trekking ability.
- Transmitted by tse-tse flies (*Glossina morsitans* group in open savanna habitats and *Glossina palpalis* group in riverine habitats); in adjacent areas without tse-tse flies, transmission of Trypanosomes by biting flies can play a role (especially for *Trypanosoma vivax*).
- Present in all sub-humid regions (“Sudan Savanna”) of West Africa where tse-tse flies occur.
- The negative effect on productivity of sheep populations is lower than in cattle because sheep are bitten by tse-tse less frequently than bovines; there is almost no impact on productivity of goats, which are only very sporadically bitten (and infected) by tse-tse flies.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Increased humidity favors increased tree cover (long-term) and enhances range, reproduction, and feeding activity of tse-tse with more frequent trypanosome infection of sheep
- Increased aridity can lead to emergence of trypanosome transmission by biting flies in areas that become unsuitable for tse-tse
- Under humid scenario shift towards trypano-tolerant sheep and goats
- In an arid CC scenario possibility of new localized tse-tse habitats in expanding irrigation areas

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Shrinking range of disease and its vector, only present in southern parts of Mali	Hot/Wet		<b>Higher losses</b> because of expanded range of tse-tse vector; lower impact than in cattle	Change to more trypano-tolerant sheep breeds or to goats instead of sheep
	Hot/Dry	<b>No losses;</b> tse-tse vector and disease may disappear		

## ENVIRONMENTALLY TRANSMITTED DISEASES

### ANTHRAX<sup>xxxiii</sup>

- Spore-forming bacteria (*Bacillus anthracis*) found in soil and particularly in sediment of ponds and lakes; spores survive in the environment for decades, if not for centuries; when ingested during grazing or drinking the spores germinate into bacteria and cause a peracute septicemia that leads to instant death in sheep; important zoonosis.
- When carcasses are opened (e.g., by scavengers), bacteria form spores, which again contaminate the environment; spores can be carried by floods onto clean pastures.
- Sporadic outbreaks occur particularly in dry season when sheep graze in dry flood zones and gather at overcrowded contaminated watering ponds; the disease is extremely aggressive and brief in sheep, which die suddenly in large numbers without showing clinical symptoms.

#### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario increased availability of grazing throughout the year with sheep and goats spending shorter time grazing in risky flood zones
- Increased aridity would see sheep and goat herds spend longer time in risky areas
- Increased flooding can spread infection onto new pastures, possible under both CC scenarios (sporadic flash floods also occur with arid CC)

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic on certain pastures, esp. in flood zones	Hot/Wet	Less time spent grazing on infected pastures → <b>Lower losses</b>		
	Hot/Dry		Longer time spent grazing on infected pastures → <b>Higher losses</b>	Improved vaccination cover

## CLOSTRIDIOSIS<sup>xxxiv</sup>

- Spore-forming anaerobic bacteria (*Clostridium spp.*) found in soil and in intestinal flora of mammals; spores survive in the environment and are regularly ingested during grazing; a sudden change in diet allows clostridia in the guts to mass-multiply and produce lethal toxins that are rapidly absorbed from the intestine and kill especially young sheep/goats during their first grazing season within a few hours (sometimes animals are found dead); older animals develop good immunity.
- Following good rainfall with fast growth of young lush grass the sudden change in diet destabilizes the intestinal conditions and triggers mass-multiplication and toxin production of clostridia; this does not represent a true infection because the pathogens are already part of the normal intestinal flora (= enterotoxaemia).
- Occurrence of pathogenic clostridia in pastures is not uniform, but is affected by soil type (content of organic matter, pH); sporadic seasonal outbreaks occur more often in certain areas than in others; immunity of adult sheep/goats depends on previous exposure and cannot always be relied upon.
- Economic importance because large numbers of young sheep/goats are lost suddenly.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario more frequent grazing of sheep on young lush growth, increases likelihood of outbreaks
- With humid CC scenario sheep and goats could also move further north and infest currently arid pastures with clostridia spores

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in sheep worldwide, occurs in soils rich in organic substance, seasonal outbreaks	Hot/Wet		Disease occurs more frequently → <b>Higher losses</b>	Improved vaccination cover
	Hot/Dry	Disease occurs less frequently → <b>Lower losses</b>		

## FLUKE (FASCIOLA)<sup>xxxv</sup>

- Trematode parasites (in the Sahel, esp. *Fasciola gigantica*), adult stages parasitize the bile ducts of the liver in ruminants. The infective larval stage causes tissue damage due to mass migration through the tissue of the liver and other internal organs, resulting in untreatable massive internal bleeding and sudden death especially in young sheep. The adult stage causes chronic loss of red blood cells and blood plasma resulting in persistent wasting disease in older sheep; this effect much more severe and progresses to a critical stage much faster than it does in cattle.
- Indirect transmission cycle includes environmental phase with developmental parasite stages living in water snails (= intermediate hosts); released infectious larvae encyst on vegetation and re-enter sheep host during grazing on fluke-infested pasture.
- Occurs in localized aquatic habitats (near permanent water and in flood zones), including in irrigation schemes (*Fasciola gigantica* gains new habitats through construction of water reservoirs and dams).
- Acute fasciolosis can lead to significant losses of young sheep and severe loss in condition in adult sheep; unless strategic prophylactic treatments are applied, keeping sheep in fluke infested areas is uneconomical.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario creates more suitable snail habitats that support parasite transmission
- Increased aridity reduces snail habitats
- Under humid CC scenario, herds depend less on fluke infested grazing; with increased aridity herds are more likely to graze on fluke infested pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in specific habitats (e.g. around water reservoirs, irrigation areas)	Hot/Wet	Less time spent grazing risky pasture → <b>Lower losses</b>		
	Hot/Dry		Longer time spent grazing risky pastures → <b>Higher losses</b> ; higher impact than in cattle	Strategic anti-parasite treatment before losses occur

## FOOTROT<sup>xxxvi</sup>

- Bacterial pathogens (*Dichelobacter nodosus*) cause footrot in sheep and occasionally in goats, manifesting as severe lameness with sheep being unable to walk and graze. Healing of the hooves is very protracted and often incomplete. Susceptibility is breed-dependent.
- Transmission does not require direct contact and occurs mostly via contaminated environment (pasture, trekking routes); the duration of survival of the pathogen in the environment is closely controlled by climatic conditions (humidity, ultraviolet radiation); unprotected watering points important for transmission under arid conditions.
- Causes severe drop in body condition and productivity as well as the need to cull many sheep that remain with malformed hooves.
- Footrot is common and economically very important in most sheep-keeping regions; in semi-arid regions, brief seasonal outbreaks can affect large numbers of sheep.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid pastures are an absolute precondition for pathogen transmission & disease
- Management can mitigate humid CC effect on ranches and in closed farms, but this is very difficult under communal grazing

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in sheep worldwide, strictly seasonal outbreaks linked to wet conditions	Hot/Wet		Disease occurs much more frequently → significantly <b>increased production losses</b>	Improved vaccination cover, foot baths
	Hot/Dry	Disease absent over long periods → significantly <b>reduced production losses</b>		

## GASTRO-INTESTINAL HELMINTHS<sup>xxxvii</sup>

- Gastro-intestinal parasites, adult stages live in the stomach or intestine. Some species attach to the lining of the stomach or the intestine and suck blood. Pathogenicity varies depending on the level of infestation and on helminth species. They interfere with digestion and some can cause severe anemia (in sheep especially *Haemonchus contortus*).
- Transmission cycle includes environmental larval phase that lives on pasture vegetation. Infective larvae are only active on pasture under humid conditions, and survive only for a rather limited period on pastures under hot and arid conditions; transmission in the semi-arid areas occurs seasonally during rains.
- High infection rate causes wasting disease and interferes massively with lamb growth (deaths possible); adult sheep and goats do not develop resistance and are severely affected by high burdens of gastro-intestinal helminthes. This is the single most important disease factor affecting productivity of sheep and goat flocks worldwide – but is of less economic importance under extensive highly mobile management in arid regions.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario lifespan and activity of infective larvae on pasture increases, resulting in more efficient transmission
- Increased aridity reduces efficiency of transmission
- Herd mobility is a modulating factor; humid CC impact is felt more in stationary farm-based animals grazing on permanent pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in all camel keeping regions	Hot/Wet		Heavier endoparasite burden → <b>Increased production losses</b> ; much higher impact than in cattle – likely to gain <b>major importance</b> in sheep/goats	Anti-parasite treatment timed in relation to expected GI-Helminth risk; breed helminth resistant sheep
	Hot/Dry	Lighter endoparasite burden → <b>Reduced production losses</b>		

## SALMONELLOSIS<sup>xxxviii</sup>

- Bacteria living in the gastro-intestinal tract of humans and animals; more than 2000 salmonella serotypes (including a sheep-specific type *Salmonella abortus ovis*) cause different clinical diseases incl. septicemia, diarrhea and abortions; affect mainly young animals; important zoonosis.
- Fecal-oral transmission via diseased and healthy salmonella carriers that contaminate the environment with their feces; salmonella can survive for long periods in water and in wet, shaded, micro-environments.
- Salmonella is more prevalent under intensive husbandry conditions; transmission in extensive production systems in the semi-arid areas occurs mainly via crowded and contaminated night enclosures.
- Lack of stable endemic immunity, and common presence of healthy carriers, leads to persistent losses through abortions and from deaths in young animals. Cumulative production losses are often overlooked.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario results in enhanced pathogen survival and multiplication of pathogens in micro-environments leading to a higher salmonella infection rate in sheep and goats
- Increased aridity reduces exposure of animals to salmonella
- Transmission via contaminated overnight enclosures in mobile pastoralist systems is more affected by increased aridity than transmission in housed farm animals grazing on permanent pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet		Enhanced pathogen survival in micro-environments → Slightly <b>increased losses</b>	Improved vaccination cover
	Hot/Dry	Slightly <b>reduced losses</b> esp. under mobile management system		

## ANIMAL-ANIMAL TRANSMITTED DISEASES

### CONTAGIOUS CAPRINE PLEURO-PNEUMONIA (CCPP)<sup>xxxix</sup>

- Bacterial pathogen (*Mycoplasma capricolum ssp. capripneumoniae*, formerly referred to as “*Mycoplasma* strain F38”) causes a severe respiratory disease in goats with 100 percent morbidity and 80 percent mortality upon exposure of non-immune naïve herds; surviving goats develop chronic progressive pneumonia that does not respond well to antibiotic treatment and are often unable to maintain sufficient lactation for raising their kids.
- Transmission takes place via aerosols containing infectious droplets and requires close contact; does not infect sheep.
- Unless controlled, CCPP is common under intensive and extensive husbandry conditions; affected animals and whole goat herds become unproductive and have to be slaughtered.
- CCPP interferes with pastoralist milk production and severely affects cash income through loss in market value and reduced sales of goats.

#### Climate change influence on pathogen-transmission, livestock management, land use

- Humid conditions favor pathogen survival in infectious aerosols with more efficient transmission
- Arid conditions with poor nutrition lower the resistance of goats to this infection
- Under arid CC scenario higher risk of pathogen transmission due to more frequent herd contacts at fewer watering points and on overstocked pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet	<b>No change</b>		Improved vaccination cover
	Hot/Dry	<b>No change</b> ; more frequent outbreaks possible in highly mobile pastoralist livestock		Improved movement control and systematic vaccination campaign

## ORF (PUSTULAR DERMATITIS)<sup>x1</sup>

- Viral pathogen (*Parapoxvirus ovis*) endemic in all sheep-keeping regions; pox-like lesions form mostly on the head, lips, and mouth in young suckling lambs and kids. Healing is without problems unless lactating ewes / goats are underfed and give little milk → lack of milk results in weak lambs/kids and potentially high losses.
- Transmitted by direct contact; the Orf virus of sheep, goats, and camels may be identical, and infection can be transmitted and shared between the three livestock species.
- Orf occurs regularly in relation to lambing rhythm of the herd.
- Economic importance because of loss or stunting of lambs and kids.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under wet unhygienic conditions secondary bacterial infections complicate healing of Orf lesions resulting in prolonged and more severe disease
- In arid situations, there can be poor nutritional status and low milk yield of ewes and goats with weaker lambs and kids that suffer higher mortality

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali and all sheep keeping regions	Hot/Wet		More severe clinical disease → <b>higher losses</b>	Improved vaccination cover
	Hot/Dry	Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)		

## OVINE BRUCELLOSIS<sup>xli</sup>

- Bacterial pathogen (*Brucella melitensis*) causing abortion in sheep and goats and also leading to persistent healthy adult carriers; it is excreted in the milk of healthy carriers (=persistently infected adult goats and sheep) – important zoonosis in areas with common raw milk consumption.
- Transmission via the environment (stables, enclosures, paddocks) which is contaminated with *Brucella* during abortion; this route plays a by far greater role than occasional venereal transmission; *Brucella* can survive for a short period in humid shaded micro-environments.
- Unless controlled, *Brucella* is common under intensive and extensive husbandry conditions; transmission in extensive production systems in the semi-arid areas occurs mainly via contaminated night enclosures.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid scenario results in better pathogen survival on pasture and more efficient transmission
- Increased aridity reduces transmission on pasture
- Transmission via contaminated pastures and overnight enclosures in mobile pastoralist systems is more affected by increased aridity than transmission in housed farm animals on permanent pastures

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet	<b>No change</b>		Improved vaccination cover
	Hot/Dry	<b>No change</b>		

## PESTE DES PETITS RUMINANTS (PPR) (A.K.A. GOAT PLAGUE)<sup>xliii</sup>

- Viral pathogen endemic to West Africa, causing acute pneumo-enteritis (respiratory tract and intestine affected → diarrhea and pneumonia) in goats and sheep; more severe in goats than in sheep; in naïve herds infected for the first time, morbidity in goats is 100 percent and mortality 90 percent; potentially high losses also occur in sheep (40-60 percent mortality). In endemic situation losses are much lower, affecting especially young animals.
- Historically more common in sub-humid and humid parts of West-Africa, now occurs in all parts of the Sahel from Atlantic to the Indian ocean and in Asia.
- Transmitted by close direct contact; the virus is very sensitive to heat and to sunlight, and it does not survive outside the host.
- The most deadly and economically the most important infectious disease and the most important single cause of disease and death in small ruminants (especially goats); currently targeted by OIE/FAO for worldwide eradication (similar to Rinderpest).
- Poses a serious threat to livestock dependent pastoralist and agro-pastoralist livelihoods, especially where PPR epidemics affect non-endemic regions.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under arid conditions transmission becomes more likely due to much more frequent herd contacts at watering points and on overstocked pastures; an arid scenario also favors introduction into new areas via unusual long distance herd migration
- Humid conditions and shade (lower temperature) favor transmission of virus

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
PPR range covers the whole Sahel – including Mali; endemic with sporadic outbreaks, Africa-wide eradication efforts about to begin	Hot/Wet	<b>No change</b>		Worldwide eradication of PPR foreseen → disease will hopefully disappear from the Sahel and Africa in the future
	Hot/Dry	<b>No change</b> ; more frequent outbreaks possible in highly mobile pastoralist livestock		

## SHEEP & GOAT POX<sup>xliii</sup>

- Viral pathogen causing fever, abortions, typical pox lesions on the skin and lesions on the inner surfaces of the respiratory tract; self-limiting, but secondary respiratory infections can cause multiple deaths, especially in weak lambs and kids (this is directly related to nutritional status and milk yield of lactating ewes/goats).
- Transmitted by direct contact; under humid conditions biting insects transmit bacteria that super-infect skin lesions, which significantly prolongs the healing process and weakens lambs/kids severely.
- Occurs in periodic epidemics that depend largely on the lambing cycle and on the immune status of the sheep/goat herd.
- Economic importance is due to loss and stunting in young lambs/kids.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid conditions flies act as mechanical transmitters of bacteria that super-infect pox lesions and secondary respiratory infections can kill lambs/kids
- In arid situation there can be poor nutritional status and low milk yield of ewes/does with weaker lambs/kids that suffer more from pox
- With arid CC scenario there can be more frequent herd contact at watering points and enhanced spread of more virulent strains

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in Mali	Hot/Wet		More severe clinical disease → <b>higher losses</b>	Improved vaccination cover
	Hot/Dry	Less severe clinical disease → <b>lower losses</b> (modulating effect of nutritional status)		

# CHICKENS

While trade in chickens, in one-day-chicks, and in eggs plays a major role in pathogen transmission the distinction between vector-borne, environmental, and animal-to-animal transmission is not always as clear cut in chickens as it is in other livestock.

## VECTOR-BORNE DISEASES

### FOWL POX<sup>xliv</sup>

- A chicken-specific pox virus causing typical pox lesions and also generalized disease in chicken of all age groups; protracted outbreaks; generalized form causes mortalities.
- The infection is transmitted by mosquitoes (*Culex spp.*); once infected, these harbor infectious virus for months – disease prevalence is linked to seasonal vector density and activity; virus is also spread by direct contact.
- Stress leads to flare-up of latent pox infection.
- Important mainly because of production losses over extended periods.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Heat stress → flare-up of latent pox infection;
- Higher rainfall → enhanced mosquito reproduction & activity → increase in virus transmission;
- Increased aridity → reduced mosquito population (e.g. *Culex quinquefasciatus*) → less frequent and less efficient pox transmission
- Increased aridity → reduced availability and increased prices of chicken feed → chicken in poor body condition and more susceptible

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Present in Mali	Hot/Wet		Increased risk of outbreaks and <b>higher production losses</b>	Improved vaccination cover
	Hot/Dry	Reduced risk of outbreaks → <b>lower production losses</b>		Ensure adequate quantity and quality of chicken feed

## ENVIRONMENTALLY TRANSMITTED DISEASES

### AVIAN COCCIDIOSIS<sup>xlv</sup>

- Decreased growth rate, severe diarrhea, and high mortality (depends on infection level) caused by 51 coccidia (*Eimeria spp.*), intestinal protozoan parasites that are universally present in chicken flocks. Coccidia can survive for very long periods in contaminated environment depending on environmental conditions (51 coccidia are resistant to disinfection). Disease only occurs after ingestion of large numbers of coccidia.
- Important because it cannot be eradicated from chicken flocks; requires frequent medication, poses permanent mortality threat especially to younger birds, and causes constant production losses.

#### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario there is a longer survival and higher buildup of coccidia in the environment
- Increased aridity → reduced availability and increased prices of chicken feed → chicken in poor body condition and more susceptible

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in all chicken-keeping regions	Hot/Wet		More frequent disease → <b>higher losses</b>	Prophylactic anti-coccidial drugs
	Hot/Dry	Less frequent disease → <b>lower losses</b> (modulating effect of poor body condition in relation to affordability/availability of feed)		Ensure adequate quantity and quality of chicken feed

## GASTRO-INTESTINAL HELMINTHS<sup>xlvi</sup>

- Gastro-intestinal parasites, helminths are especially important in free ranging chickens; adult stages live in the gizzard, trachea, esophagus, and small and large intestine; pathogenicity varies considerably depending on the level of infestation and on helminth species; they interfere with digestion and some can cause anemia and death.
- Transmission cycle includes environmental larval phase, many of which require an intermediate host for their life-cycle (ants, cockroaches, snails, earthworms, beetles, grasshoppers etc.); life-cycles to produce infective larvae are sensitive to hot and arid conditions.
- High infection rates cause wasting disease and even death → potentially high and continuous negative impact on production.

### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario lifespan and activity of infective larvae and intermediate hosts increases → more efficient transmission
- Increased aridity reduces efficiency of transmission → reduced exposure

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in free ranging chickens worldwide	Hot/Wet		Heavier endoparasite burden → <b>increased losses</b>	Anti-parasite treatment
	Hot/Dry	Lighter endoparasite burden → <b>reduced losses</b>		

## GUMBORO DISEASE<sup>xlvii</sup>

- Viral pathogen is endemic in all chicken-keeping regions; it causes severe long lasting immune-suppression (subclinical form) or weakness, in-coordination, and diarrhea (acute form). Mortality can be as high as 20 percent,
- Transmitted by direct and indirect contact, very stable in the environment → difficult to eradicate from infected premises.
- Economic importance because of suppressed production and mortalities.

### Climate change influence on pathogen-transmission, livestock management, land use

- The effect of both humid and arid CC scenarios is limited because the virus survives for long periods in contaminated premises

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken-keeping regions	Hot/Wet	<b>Stable losses</b>		Improved vaccination cover
	Hot/Dry			

## SALMONELLOSIS<sup>xlviii</sup>

- Bacteria living in the gastro-intestinal tract of humans and animals; more than 2000 *Salmonella* serotypes are known worldwide (including some few host-specific types) and cause different clinical diseases incl. poor growth; weakness; diarrhea; septicemia & mortality is seen mainly in young chicken; high prevalence in chicken flocks; important zoonosis and of major food safety concern.
- Fecal-oral transmission occurs via salmonella carriers that contaminate the environment with their feces; salmonella can survive for long periods in water and in wet shaded micro-environments.
- Transmission in chickens is mainly via crowded and contaminated premises.
- Persistent losses are seen, especially in young chickens, as is contamination of chicken products entering the food chain.

### Climate change influence on pathogen-transmission, livestock management, land use

- Humid CC scenario results in enhanced pathogen survival and multiplication inside premises → higher salmonella exposure in chicken;
- Increased aridity → reduced exposure of chicken to salmonella
- Increased aridity → poor availability and increased prices of chicken feed → chickens in poor body condition are more susceptible

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken-keeping regions	Hot/Wet		<b>Higher losses</b>	Improved hygiene on premises
	Hot/Dry	<b>Reduced losses</b> (modulating effect of poor body condition in relation to affordability/availability of feed)		Ensure adequate quantity and quality of chicken feed

## ANIMAL-ANIMAL TRANSMITTED DISEASES

### NEWCASTLE DISEASE (ND)<sup>xlix</sup>

- Occurs in periodic epidemics that depend largely on the immune status of the local chicken population and/or introduction of highly virulent virus strains; produces mostly acute respiratory symptoms, but also nervous signs and diarrhea; mortality varies extremely and depends on virus strain – velogenic (= highly virulent) Newcastle disease is endemic in many parts of Africa and regularly destroys whole village chicken populations.
- Fecal-oral and aerosol transmission occur; aerosol transmission accelerated by wind results in much faster and wider spread within localities.
- Velogenic (= highly virulent) Newcastle disease is the most important infection in chicken and the single most important obstacle to improved chicken productivity in Africa.

### Climate change influence on pathogen-transmission, livestock management, land use

- Wind speed affects local transmission
- In a more arid scenario there are reduced habitats for overwintering migratory birds; these then concentrate in smaller wetlands with increased risk of spread of velogenic virus strains
- Arid and humid CC scenarios can both alter routes of migratory birds

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken-keeping regions, presence of very pathogenic (velogenic) ND virus strains	Hot/Wet	<b>No change</b>		
	Hot/Dry		More frequent Newcastle Disease outbreaks → <b>increased losses</b>	Improved vaccination cover

## HIGHLY PATHOGENIC AVIAN INFLUENZA (HPAI)<sup>1</sup>

- Occurs in worldwide periodic pandemics; clinical picture extremely variable depending on the HPAI strain; varies from acute febrile respiratory disease with very high losses to clinically almost unapparent mild infections; high zoonotic potential capable of causing worldwide human influenza pandemics – mild and acute forms are both very dangerous for humans.
- Transmission occurs through trade in live chickens and eggs, as well as being spread by wild migratory birds; migratory routes of wild birds are affected by CC; co-evolution has allowed different species of wild birds to resist and tolerate HPAI infection without impediment of their migratory range and ability.
- Direct losses related to high mortality/destruction of infected flocks, reduced egg production, and decreased weight gains.
- Indirect costs include clean-up, complete disruption of trade, disruptions in production schedules, costs for restocking, and lost income.

### Climate change influence on pathogen-transmission, livestock management, land use

- More arid climate → reduced wetland habitats for over-wintering migratory birds that concentrate in smaller areas with increased risk of spread of HPAI
- Arid and humid CC scenario can both alter routes of migratory birds
- Housing chicken reduces transmission risk; main spread of HPAI is through trade

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Presence related to concentrations of over-wintering migratory birds in wetlands; not a major problem in Mali at present	Hot/Wet	<b>No major change</b>		Raised alertness and prompt stamping out of infected flocks to prevent further spread
	Hot/Dry		<b>Increased risk</b> because of higher concentration of migratory birds in fewer over-wintering habitats	

## PULLORUM DISEASE<sup>li</sup>

- Host specific salmonella (*Salmonella pullorum*) causes very high losses (up to 100 percent) in chicks and young chicken.
- This is transmitted primarily through the egg (chick already infected at hatching); also transmitted by direct contact; shed by carriers.
- Occurs in all chicken-rearing regions where it has not been eradicated.
- Great economic importance because of high mortality.

### Climate change influence on pathogen-transmission, livestock management, land use

- Transmission via eggs not affected by CC
- Heat stress in housed chickens could potentially increase disease severity

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken-keeping regions	Hot/Wet	<b>No change</b>		Improved vaccination cover
	Hot/Dry			

## MAREK DISEASE<sup>lii</sup>

- Viral pathogen endemic in all chicken-keeping regions worldwide; strains vary widely in virulence; can cause tumors and mortality in up to 80 percent of young birds; causes transient paralysis, depression, and death.
- Easily transmitted by direct contact; virus can also survive in litter for months; stress leads to resurgence of occult infection.
- Economic importance because of mortality.

### Climate change influence on pathogen-transmission, livestock management, land use

- Transmission not affected by CC
- Heat stress can reactivate occult infections in housed chicken

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken keeping regions	Hot/Wet	Slightly <b>increased risk</b> of outbreaks (increased heat stress)		Improved vaccination cover
	Hot/Dry			

### AVIAN INFECTIOUS BRONCHITIS<sup>liii</sup>

- Viral respiratory pathogen (a Corona-virus) endemic in all chicken-keeping regions worldwide; outbreaks occur in relation to flock/population immunity; strains vary widely in virulence; some strains can cause kidney failure and high mortality.
- Easily transmitted by direct contact and infectious aerosols; stress affects severity of outbreaks.
- Economic importance because of severe loss in production while 100 percent of the flock suffers from respiratory disease.

#### Climate change influence on pathogen-transmission, livestock management, land use

- Transmission not affected by CC; heat stress exacerbates clinical disease in housed chickens

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken-keeping regions	Hot/Wet	More severe outbreaks (heat stress) → <b>higher production losses</b>		Improved vaccination cover
	Hot/Dry			

### AVIAN MYCOPLASMOSIS<sup>liv</sup>

- Bacterial respiratory pathogen (esp. *Mycoplasma gallisepticum*) endemic in all chicken-keeping regions worldwide; causes chronic respiratory disease in chicken flocks; strains vary in virulence.
- Transmission occurs via infected eggs, but it is also easily transmitted by infectious aerosols; stress affects severity of outbreaks.
- Economic importance because of severe loss in egg production and frequent condemnation of carcasses of birds that have recovered from respiratory disease.

#### Climate change influence on pathogen-transmission, livestock management, land use

- Heat stress exacerbates clinical disease in housed chickens

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chickenskeeping regions	Hot/Wet	More severe outbreaks (heat stress) → <b>higher production losses</b>		Improved vaccination cover
	Hot/Dry			

## FOWL CHOLERA<sup>iv</sup>

- Various bird-specific *Pasteurella multocida* serotypes cause sporadic sudden septicemia (“sudden death”) with very high losses (up to 100 percent) in chicken; survivors become chronic carriers.
- Transmitted primarily through direct contact; shed by carriers.
- Occurs in all chicken-rearing regions.
- Of economic importance because of sudden high mortality.

### Climate change influence on pathogen-transmission, livestock management, land use

- Outbreaks triggered by introduction of virulent *Pasteurella* serotypes into unprotected flocks; no CC impact; trade is a major risk

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Endemic in chicken keeping regions	Hot/Wet	<b>No change</b>		Improved vaccination cover (difficult due to multiplicity of serotypes)
	Hot/Dry			

# DONKEYS, MULES, AND HORSES

## VECTOR-BORNE DISEASES

### AFRICAN HORSE SICKNESS<sup>lv</sup>

- Arthropod-borne viral pathogen (an “Arbovirus”) affecting particularly horses (mortality 70 percent to 95 percent) – but also mules (mortality 50 percent to 70 percent) and donkeys (different breeds possess different levels of resistance; very mild in local donkeys in endemic areas); it causes severe inflammation of the lungs and severe respiratory distress (peracute pulmonary form → death within hours) or swelling of the head and neck and stiffness (death after three to six days). Extreme variability is seen in clinical manifestation (mild in endemic areas in animals in good body condition).
- It is transmitted by midges (*Culicoides spp.*), which can be wind-carried over long distances; the virus is remarkably heat-stable and survives in tissues of dead animals for very long periods; disease in endemic areas is seasonal directly related to vector activity. AHS episodes outside endemic areas have occurred in relation to movement of infected horses/donkeys, with high rainfall and strong winds causing high mortalities in naïve horses, mules, and donkeys in North Africa and also in Spain.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Higher rainfall enhances reproduction and biting activity of midges and increases virus transmission
- Wind speed affects long distance spread of infected midges
- Concentration of donkeys/mules in irrigation areas offers favorable conditions for midges and for disease transmission

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Not reported from Mali, present in neighboring countries	Hot/Wet		Increased risk of disease introduction	Movement control at borders for disease free countries; improved vaccination cover in infected countries
	Hot/Dry	Reduced risk of disease introduction		

## TRYPANOSOMOSIS (INCL. TRYPANOSMA EVANSI)<sup>lvii</sup>

- The tse-tse transmitted *Trypanosoma vivax* and *Trypanosoma brucei*, and to a lesser extent also *Trypanosoma congolense*, infect horses and donkeys and cause intermittent fever, lasting immune-suppression, progressive anemia, progressive weight loss, and eventually death. In horses and donkeys *Trypanosoma brucei* can cause death within one to three weeks; the Tabanid transmitted *Trypanosma evansi* causes a chronic disease in horses similar to the one observed in camels. Chronic Trypanosomosis cases do not respond well to treatment; they cannot be used for traction or for any other work, and are often culled.
- Transmitted by tse-tse flies (*Glossina morsitans* group in open savanna habitats and *Glossina palpalis* group in riverine habitats) and by biting flies (Tabanids); frequency of transmission and disease is directly related to seasonal increase in vector population / biting activity.
- Present in all sub-humid and semi-arid parts of the Sahel.
- Can pose a threat to crop production because increased vector activity overlaps with the planting and growing season.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Increased humidity enhances range, reproduction and feeding activity of tse-tse and other biting flies (*Tabanus*, *Stomoxys*) → donkeys will be more often infected with trypanosomes
- Under arid scenario and/or more intensive land cultivation *Glossina morsitans* is likely to disappear

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Tse-tse transmitted trypanosomes only in southern Mali (shrinking range); <i>T. evansi</i> present country-wide	Hot/Wet		Higher transmission risk in wider area → <b>Increased loss of traction power</b>	Improved tse-tse control (traps) and use of repellents
	Hot/Dry	Lower transmission risk for <i>T. evansi</i> ; tse-tse transmitted trypanosomes may disappear → <b>Reduced loss of traction power</b>		

## ENVIRONMENTALLY TRANSMITTED DISEASES

### GASTRO-INTESTINAL HELMINTHS AND LIVER FLUKE<sup>lviii</sup>

- Can affect horses in a similar way as other herbivorous livestock. Lung-worms transmitted via pasture can play an important role in donkeys, but require cool highland climate for efficient transmission.

#### Climate change influence on pathogen-transmission, livestock management, land use

- Under humid CC scenario lifespan, activity and transmission of flukes and helminths increases

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
GI-helminths present country-wide, flukes only in specific habitats (water reservoirs, ponds)	Hot/Wet		Higher exposure → <b>Increased loss of traction power</b>	Anti-parasite treatment against helminths/flukes
	Hot/Dry	Lower exposure → <b>Reduced loss of traction power</b>		

## GLANDERS<sup>lix</sup>

- Bacterial pathogen causes severe lung and also skin disease in horses and donkeys. Donkeys are more susceptible than horses and more likely to develop the acute form with high mortality, while horses develop a more chronic form. Horses may recover, but there is no stable immunity (depends on body condition).
- Transmission takes place mainly via contaminated environment (pathogen survives in water for up to 20 days) and through very close contact.
- Movement of sub-clinically infected horses is the most important means of spreading the infection.
- Dangerous zoonosis.

### Climate change influence on pathogen-transmission, livestock management, land use

- Effect of humid and arid CC scenario both limited because pathogen is mainly spread by sub-clinically infected carriers
- Donkeys are more important for plowing and are also much more susceptible than horses

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Absent from Mali, but reported from neighboring Mauritania	Hot/Wet Hot/Dry	<b>No CC related change</b> , veterinary control measures can prevent introduction into Mali		Improved cross-border movement control for imported horses to prevent spread to other countries in the Sahel

# PIGS

## VECTOR-BORNE DISEASES

### AFRICAN SWINE FEVER (ASF)<sup>x</sup>

- Tick-borne viral pathogen affects domestic pigs, causing mostly acute disease (mortality up to 100 percent within two to nine days). Local African domestic pig breeds are more resistant, and chronically infected survivors can occur in endemic areas. ASF causes mass abortions and death.
- A natural reservoir exists in wild warthogs that do not develop overt disease (the role of bushpigs as reservoir is less clear); it is transmitted between warthogs and free ranging domestic pigs by soft ticks (*Ornithodoros spp.*). Transmission between domestic pigs is entirely by direct contact; once the infection is established in domestic pigs the tick vector no longer plays a role in transmission; the virus survives only three days in pig houses. In West Africa ASF has not been found in warthogs, but outbreaks in domestic pigs do occur at intervals. ASF has spread outside endemic areas to Southern Europe and to Russia
- This is the most deadly infection in domestic pigs.

### Climate change influence on vector, pathogen-transmission, livestock management, land use

- Higher rainfall → enhanced reproduction and activity of vector ticks → increased risk of virus transmission between wild warthogs and free ranging domestic pigs
- Adequate housing fully disrupts transmission between reservoir in the wild and domestic pigs

Current Disease Status in Mali	Climate Change Scenario	Expected Climate Change Impacts by 2025		Possible mitigation strategy
		Disease unlikely to gain in importance	Disease likely to gain in importance	
Not a major problem in Mali at present	Hot/Wet		<b>Increased risk of transmission in free ranging domestic pigs</b>	Development of an ASF vaccine (research currently ongoing)
	Hot/Dry	Reduced risk of transmission and <b>reduced importance</b> of pig industry (rising feed costs)		

Note on sources: In addition to the texts listed in the endnotes, the author relied upon the following websites of the World Organisation for Animal Health (OIE) for additional information (all OIE websites accessed August, 2013): OIE World Animal Health Information Database (WAHID), [http://www.oie.int/wahis\\_2/public/wahid.php/Wahidhome/Home](http://www.oie.int/wahis_2/public/wahid.php/Wahidhome/Home); and OIE WAHID Country Timelines, [http://www.oie.int/wahis\\_2/public/wahid.php/Countryinformation/Countrytimelines](http://www.oie.int/wahis_2/public/wahid.php/Countryinformation/Countrytimelines).

---

- <sup>i</sup> Potgieter, F.T., & Stoltsz, W.H. Bovine anaplasmosis. (1994). Chapter 41 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>ii</sup> De Vos, A.J., & Potgieter, F.T. (1994). Bovine babesiosis. Chapter 23 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>iii</sup> Barnard, B.J.H., Munz, E., Dumbell, K., & Prozesky, L. (1994). Lumpy skin disease. Chapter 52 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>iv</sup> Swanepoel, R.S., & Coetzer, J.A.W. (1994). Rift Valley Fever. Chapter 71 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>v</sup> Pipano, E. (1994). Theileria annulata theileriosis. Chapter 34 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>vi</sup> Brown, C.D.G., Hunter, A.G., & Luckins, A.G. (1992). Diseases caused by Protozoa – Tropical Theileriosis, pp. 191-196 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), *Handbook on Animal Disease in the Tropics*. British Veterinary Association & Bailliere Tindall, ELBS 4<sup>th</sup> edition, London, UK.
- <sup>vii</sup> Connor, R.J. African animal trypanosomiases. (1994). Chapter 11 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>viii</sup> De Vos, V. (1994). Anthrax. Chapter 153 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>ix</sup> Hammond, J.A., & Sewell, M.M.H. (1992). Diseases caused by helminths – Fasciolosis, pp. 119-123 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), *Handbook on Animal Disease in the Tropics*. British Veterinary Association & Bailliere Tindall, ELBS 4<sup>th</sup> edition, London, UK.
- <sup>x</sup> Hammond, J.A., & Sewell, M.M.H. (1992). Diseases caused by helminths – Trichostrongylidosis, pp. 149-155 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), *Handbook on Animal Disease in the Tropics*. British Veterinary Association and Bailliere Tindall, ELBS 4<sup>th</sup> edition, London, UK.
- <sup>xi</sup> Venter, B.J., Myburgh, J.G., & van der Walt, M.L. (1994). Bovine salmonellosis. Chapter 130 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xii</sup> Bishop, G.C., Bosman, P.P., & Herr, S. (1994). Bovine brucellosis. Chapter 124 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xiii</sup> Schneider, H.P., van der Lugt, J.J., & Hübschle, O.J.B. (1994). Contagious bovine pleuropneumonia. Chapter 177 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xiv</sup> Van Tonder, E.M., & Horner, R.F. (1994). Dermatophilosis. Chapter 176 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xv</sup> Thomson, G.R. Foot and Mouth Disease. (1994). Chapter 89 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

- 
- <sup>xvi</sup> Bastianello, S.S., & Nesbit, J.W. (1994). Haemorrhagic Septicemia. Chapter 142 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xvii</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Rift Valley Fever, pp. 228-230 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xviii</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Trypanosomosis, pp. 273-281 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xix</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Trypanosomosis, pp. 273-281 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xx</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Anthrax, pp. 33-35 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxi</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Trychostrongylidosis (Gastrointestinal Worm Infection), pp. 348-353 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxii</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Salmonellosis, pp. 73-78 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxiii</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Brucellosis, pp. 108-116 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxiv</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Camelpox, pp. 176-187 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxv</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Dermatophilosis, pp. 141-148 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxvi</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Pasteurellosis, pp. 49-54 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxvii</sup> Wernery, U., & Kaaden, O.R. (Eds.). (2002). Contagious Ecthyma, pp. 187-192 in *Infectious Diseases in Camelids*, 2<sup>nd</sup> edition, Blackwell Science, Berlin-Vienna 2002.
- <sup>xxviii</sup> Hammond, J.A., & Sewell, M.M.H. (1992). Diseases caused by helminths – Fasciolosis, pp. 119-123 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), *Handbook on Animal Disease in the Tropics*. British Veterinary Association and Bailliere Tindall, ELBS 4<sup>th</sup> edition, London, UK.
- <sup>xxix</sup> Verwoerd, D.W., & Erasmus, B.J. (1994). Bluetongue. Chapter 43 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xxx</sup> Bezuidenhout, J.D., Prozesky, L., du Plessis, J.L., & van Amsted, S.R. (1994). Heartwater. Chapter 35 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xxxi</sup> Swanepoel, R.S., Coetzer, J.A.W. (1994). Rift Valley Fever. Chapter 71 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xxxii</sup> Connor, R.J. African animal trypanosomiases. (1994). Chapter 11 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xxxiii</sup> De Vos, V. (1994). Anthrax. Chapter 153 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xxxiv</sup> Kriek, N.P.J., Odendaal, M.W., & Hunter, P. (1994). Clostridium perfringens type D enterotoxaemia. Chapter 157 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.
- <sup>xxxv</sup> Hammond, J.A., & Sewell, M.M.H. (1992). Diseases caused by helminths – Fasciolosis, pp. 119-123 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), *Handbook on Animal Disease in the Tropics*. British Veterinary Association and Bailliere Tindall, ELBS 4<sup>th</sup> edition, London, UK.
- <sup>xxxvi</sup> Venter, B.J., & van Amstel, S.R. (1994). Fusobacterium necrophorum, Dichelobacter (Bacteroides) nodosus and Bacteroides spp. Infections. Chapter 145 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious*

---

*Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xxxvii</sup> Hammond, J.A., & Sewell, M.M.H. (1992). Diseases caused by helminths – Trichostrongylidosis, pp. 149-155 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), *Handbook on Animal Disease in the Tropics*. British Veterinary Association and Bailliere Tindall, ELBS 4<sup>th</sup> edition, London, UK.

<sup>xxxviii</sup> Neser, J.A. Ovine and caprine salmonellosis. (1994). Chapter 131 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xxxix</sup> Maré, C.J. (1994). Contagious Caprine Pleuro-Pneumonia (CCPP). Chapter 178 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xl</sup> Munz, E., & Dumbell, K. (1994). Orf. Chapter 54 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xli</sup> Herr, S. (1994). *Brucella melitensis* infection. Chapter 126 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xlii</sup> Rossiter, P.B., & Taylor, W.P. (1994). Peste des petits ruminants. Chapter 75 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xliii</sup> Munz, E., & Dumbell, K. (1994). Sheeppox and goatpox. Chapter 53 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>xliv</sup> Merck. (2005). Fowlpox. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2276-2278.

<sup>xlv</sup> Merck. (2005). Coccidiosis. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2201-2206.

<sup>xlvi</sup> Merck. (2005). Helminthiasis. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2231-2236.

<sup>xlvii</sup> Merck. (2005). Infectious Bursal Disease (Gumboro disease). In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2239-2240.

<sup>xlviii</sup> Merck. (2005). Salmonellosis. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2262-2264.

<sup>xlix</sup> Merck. (2005). Newcastle Disease & other Paramyxovirus infections. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2255-2257.

<sup>l</sup> Merck. (2005). Avian Influenza (Fowl plague). In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2297-2298.

<sup>li</sup> Merck. (2005). Salmonellosis. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2262-2264.

<sup>lii</sup> Merck. (2005). Neoplasms. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2248-2250.

<sup>liii</sup> Merck. (2005). Infectious Bronchitis. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2302-2304.

<sup>liv</sup> Merck. (2005). Mycoplasmosis (*Mycoplasma gallisepticum* infection). In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2242-2243.

<sup>lv</sup> Merck. (2005). Fowl Cholera. In *The Merck Veterinary Manual*. 9<sup>th</sup> ed., Merck & Co., Inc., Whitehouse Station, N.J., USA. Pp. 2228-2229.

<sup>lvi</sup> Coetzer, J.A.W., & Erasmus, B.J. African horsesickness. Chapter 44 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), (1994) *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

---

<sup>lvii</sup> Connor, R.J. African animal trypanosomiases. Chapter 11 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), (1994) *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>lviii</sup> Hammond, J.A., & Sewell, M.M.H. Diseases caused by helminths – Fasciolosis, and Diseases caused by helminths – Trichostrongylidosis , pp. 119-123 and pp. 149-155 in Sewell, M.M.H. & Brocklesby, D.W. (Eds.), (1992). *Handbook on Animal Disease in the Tropics*. British Veterinary Association and Bailliere Tindall , ELBS 4<sup>th</sup> edition , London, UK.

<sup>lix</sup> Bishop, G.C. Glanders. Chapter 122 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), (1994) *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

<sup>lx</sup> Plowright, W., & Thomson, G.R., Neser, J.A. African swine fever. Chapter 51 in Coetzer, J.A.W., Thomson, G.R. & Tustin, R.C. (Eds.), (1994) *Infectious Diseases of Livestock with special reference to Southern Africa*. Oxford University Press Southern Africa, Cape Town, South Africa, Vol. 1 & 2.

**U.S. Agency for International Development**

1300 Pennsylvania Avenue, NW

Washington, DC 20523

Tel: (202) 712-0000

Fax: (202) 216-3524

[www.usaid.gov](http://www.usaid.gov)